

An annotated checklist of the freshwater cladocerans (Crustacea: Branchiopoda: Cladocera) of Ecuador and the Galápagos Islands

**C. López^{1,2}, P.V. Mosquera³, H. Hampel^{4,5}, A.N. Neretina^{6*},
M. Alonso⁷, K. Van Damme⁸, A.A. Kotov^{6,9}**

¹ Escuela Superior Politécnica del Litoral, ESPOL, Facultad de Ciencias de la Vida y Centro del Agua y Desarrollo Sustentable, Campus Gustavo Galindo, Km. 30.5, Vía Perimetral, Guayaquil 090101, Ecuador. E-mail: cllopez@espol.edu.ec

² Departamento de Biología, Facultad Experimental de Ciencias, Universidad del Zulia, Avenida Universidad, Sector Grano de Oro, Maracaibo 4005, Venezuela.

³ Subgerencia de Gestión Ambiental de la Empresa Pública Municipal de Telecomunicaciones, Agua potable, Alcantarillado y Saneamiento (ETAPA EP), Benigno Malo y Sucre, Cuenca, Ecuador. E-mail: pmosquerav@gmail.com

⁴ Laboratorio de Ecología Acuática, Departamento de Recursos Hídricos y Ciencias Ambientales, Universidad de Cuenca, Av. 12 de Abril S/N, Cuenca, Ecuador. E-mails: hennihampel@gmail.com; henrietta.hampel@ucuenca.edu.ec

⁵ Facultad de Ciencias Químicas, Universidad de Cuenca, Av. 12 de Abril S/N, Cuenca, Ecuador.

⁶ Laboratory for Ecology of Aquatic Communities and Invasions, A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky Prospect 33, Moscow 119071, Russia. E-mail: neretina-anna2017@yandex.ru

⁷ Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona, Av. Diagonal 643, 08028 Barcelona, Spain. E-mail: malonso@ub.edu

⁸ Senckenberg Research Institute, Senckenberganlage 25, Frankfurt a.M. 60325, Germany. E-mail: kay.vandamme@gmail.com

⁹ Laboratory of Paleoclimatology, Kazan Federal University, Kremlevskaya Street 18, Kazan 420000, Russia. E-mail: alexey-a-kotov@yandex.ru

ABSTRACT: An annotated checklist of recorded species of Cladocera (Crustacea: Branchiopoda) from inland aquatic habitats in Ecuador (including the Galápagos) is provided. We revised all published records, evaluated the validity of each taxon and provided short taxonomic and biogeographical remarks for each taxon. A total of 34 valid species grouped into 23 genera was found. Presently, all attempts to analyze distributional patterns of cladocerans in Ecuador are premature due to the scarcity of adequate information. Current achievements in the inventory of cladocerans, information gaps and recommendations for future studies on cladoceran taxonomy in Ecuador are discussed.

How to cite this article: López C., Mosquera P.V., Hampel H., Neretina A.N., Alonso M., Van Damme K., Kotov A.A. 2018. An annotated checklist of the freshwater cladocerans (Crustacea: Branchiopoda: Cladocera) of Ecuador and the Galápagos Islands // *Invert. Zool.* Vol.15. No.3. P.277–291. doi: 10.15298/invertzool.15.3.06

KEY WORDS: checklist, Cladocera, biodiversity, Neotropics, South America, taxonomy.

*Corresponding author

Аннотированный список пресноводных ветвистоусых ракообразных (Crustacea: Branchiopoda: Cladocera) Эквадора и Галапагосских островов

К. Лопез^{1,2}, П.В. Москера³, Г. Хампель^{4,5}, А.Н. Неретина^{6*},
М. Алонсо⁷, К. Ван Дамме⁸, А.А. Котов^{6,9}

¹ Escuela Superior Politécnica del Litoral, ESPOL, Facultad de Ciencias de la Vida y Centro del Agua y Desarrollo Sustentable, Campus Gustavo Galindo, Km. 30.5, Vía Perimetral, Guayaquil 090101, Ecuador. E-mail: cllopez@espol.edu.ec

² Departamento de Biología, Facultad Experimental de Ciencias, Universidad del Zulia, Avenida Universidad, Sector Grano de Oro, Maracaibo 4005, Venezuela.

³ Subgerencia de Gestión Ambiental de la Empresa Pública Municipal de Telecomunicaciones, Agua potable, Alcantarillado y Saneamiento (ETAPA EP), Benigno Malo y Sucre, Cuenca, Ecuador. E-mail: pmosquerav@gmail.com

⁴ Laboratorio de Ecología Acuática, Departamento de Recursos Hídricos y Ciencias Ambientales, Universidad de Cuenca, Av. 12 de Abril S/N, Cuenca, Ecuador. E-mails: hennihampel@gmail.com; henrietta.hampel@ucuenca.edu.ec

⁵ Facultad de Ciencias Químicas, Universidad de Cuenca, Av. 12 de Abril S/N, Cuenca, Ecuador.

⁶ Лаборатория экологии водных сообществ и инвазий, Институт проблем экологии и эволюции им. А.Н. Северцова РАН, Ленинский проспект 33, Москва 119071, Россия. E-mail: neretina-anna2017@yandex.ru

⁷ Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona, Av. Diagonal 643, 08028 Barcelona, Spain. E-mail: malonso@ub.edu

⁸ Senckenberg Research Institute, Senckenberganlage 25, Frankfurt a.M. 60325, Germany. E-mail: kay.vandamme@gmail.com

⁹ Лаборатория палеоклиматологии, Казанский (Приволжский) федеральный университет, ул. Кремлевская 18, Казань 420000, Республика Татарстан, Россия. E-mail: alexey-kotov@yandex.ru

РЕЗЮМЕ: В статье представлен аннотированный список видов ветвистоусых ракообразных (Crustacea: Branchiopoda), отмеченных во внутренних водоемах Эквадора и Галапагосских островов. Мы пересмотрели все опубликованные находки, проанализировали валидность каждого таксона и привели короткие таксономические и биогеографические примечания для каждого таксона. Всего нами было выявлено 34 валидных вида кладоцер, принадлежащих к 23 родам. В настоящее время, любые попытки анализа паттернов распределения ветвистоусых ракообразных в Эквадоре являются преждевременными из-за нехватки надежной информации. Также в статье обсуждены текущие достижения в области инвентаризации кладоцер, информационные пробелы и приведены рекомендации для будущих исследований по систематике кладоцер Эквадора.

Как цитировать эту статью: López C., Mosquera P.V., Hampel H., Neretina A.N., Alonso M., Van Damme K., Kotov A.A. 2018. An annotated checklist of the freshwater cladocerans (Crustacea: Branchiopoda: Cladocera) of Ecuador and the Galápagos Islands // *Invert. Zool.* Vol.15. No.3. P.277–291. doi: 10.15298/invertzool.15.3.06

КЛЮЧЕВЫЕ СЛОВА: контрольный список, Cladocera, биоразнообразие, Неотропики, Южная Америка, систематика.

*Автор для переписки

Introduction

Investigations of the cladocerans (Crustacea: Branchiopoda: Cladocera) of the inland freshwater bodies of South America began at the end of the 19th century and were initially carried out by European taxonomists (e.g. Ihering, 1895; Sars, 1901; Stingelin, 1913; Delachaux, 1919; Brehm, 1956). Studies of the freshwater cladocerans of Ecuador began much later. Only at the beginning of the 1980s, some pioneer surveys were carried out as a part of limnological studies on the Galápagos Islands and Andean lakes (e.g. Colinviaux, Steinitz-Kannan, 1980). Since the 1980s only a handful of papers recorded the cladocerans from the inland waters of Ecuador (Gunkel, 2000; Casallas, Gunkel, 2001; Torres, Rylander, 2006; Obando, 2009; Briones, 2012; López-Blanco, Sinev, 2016; Van Colen *et al.*, 2017; Alonso *et al.*, 2017; Alonso, Kotov, 2017; Alonso, Sinev, 2017) have appeared. Compared to other countries in the tropical zone of South America (Valdivia Villar, 1988; Elmoor-Loureiro, 1998; 2000; Zoppi de Roa, López, 2008; Kotov, Fuentes-Reinés, 2015), the taxonomic knowledge on the cladocerans of Ecuador is limited. Such studies are now just beginning.

National and regional checklists are crucial for a basic knowledge on the biodiversity and such efforts allow a firm taxonomical basis for derived studies. Recently, several comprehensive checklists with taxonomical notes have been compiled for different countries where cladocerans are under intense study, e.g. India (Chatterjee *et al.*, 2013), China (Korovchinsky, 2013; Ji *et al.*, 2015; Xiang *et al.*, 2015) and Korea (Jeong *et al.*, 2014) in Asia; Colombia (Kotov, Fuentes-Reinés, 2015), Chile (De los Ríos Escalante, Kotov, 2015) and Brazil (Elmoor-Loureiro, 1998, 2018) in South America. At the same time, large revisions and checklists in Cladocera genera have helped to tackle taxonomical challenges (e.g. Van Damme *et al.*, 2010).

The aims of the present paper are to provide a formal list of the Ecuadorian (including the Galápagos Archipelago) cladocerans based on

all available literature, to evaluate the validity of all recorded species and to discuss some taxonomic and biogeographical issues. Moreover, we identify information gaps and provide recommendations for future investigations of taxonomy and biogeography of the cladocerans of Ecuadorian inland waters.

Material and methods

The list of the cladocerans of Ecuador compiled herein is based on data from literature, a revision of all published papers we could find to date. Species listed for Ecuador are discussed with special attention to their taxonomic status. Current valid species names and combinations are based on the FADA checklist (Kotov *et al.*, 2013), with some exclusions specified below. The species are attributed to three geographic and physiographic provinces of the continental territory of this country: Andean Region, Coastal Region, Amazonian Region according to Steere (1950) and to the Galápagos Islands (Fig. 1).

Results

In total, 34 valid species of Cladocera have been reported in literature from inland water bodies of Ecuador. These species belong to seven families and 23 genera. The family Chydoridae, as expected, includes the largest number of genera (13 genera) followed by the Daphniidae (3 genera). The remaining families each contain two or one genera. Chydoridae includes 15 species, followed by Daphniidae (12 species), Macrothricidae (2 species), Ilyocryptidae (2 species), Sididae (2 species), Bosminidae (2 species) and Moinidae (1 species).

Order Ctenopoda Sars, 1865

Family Sididae Baird, 1850

(1) *Diaphanosoma brachyurum* (Liévin, 1848)

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).



Fig. 1. Map of Ecuador showing main geographical regions, number of recorded species and number of sampled locations in each region.

Рис. 1. Карта Эквадора с обозначенными основными биогеографическими районами, число выявленных видов и число обследованных водоемов приведено для каждого региона.

Taxonomical remarks. *D. brachyurum* has been originally described from Europe and constitutes a complex of cryptic species with several related taxa in the Nearctic and Neotropical regions (Korovchinsky, 1992, 2004). Often, existing reports of this taxon refers to other species, even in Europe (Korovchinsky, 2000). In the Neotropics, this name is used incorrectly because this species is absent in South America (Korovchinsky, 2004). The Ecuadorian popula-

tions need to be revised and all Neotropical records of this species must be critically evaluated. As a result, this common and diverse tropical species group remains unresolved in Ecuador.

(2) *Sarsilatona serricauda* Sars, 1901

Records. Coastal Region: Lake Tembladera ($3^{\circ}29'29''\text{S}$ $79^{\circ}59'40''\text{W}$), 17 m a.s.l., Province of El Oro, identified on remains in lake sediments (López-Blanco, Sinev, 2016).

Taxonomical remarks. *S. serricauda* was described from Brazil (see Korovchinsky, 2004). It is distributed from the southern U.S.A. to northern Argentina (Korovchinsky, 2004; Kotov, Fuentes-Reinés, 2015).

Order Anomopoda Sars, 1865

Family Daphniidae Straus, 1820

(1) *Ceriodaphnia* cf. *cornuta* Sars, 1885

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. This taxon was initially described from Australia (Sars, 1885) and then found in other tropical regions. Now it is considered as a complex of closely related species, specific for each continent (Sharma, Kotov, 2013). According to DNA-barcoding data, the *C. cornuta* species complex harbors cryptic species in tropical regions of the Neotropics as well (Elías-Gutiérrez *et al.*, 2008). *C. cornuta*, sometimes listed under the name of its African congener, *C. rigaudi* Richard 1894, is commonly recorded in the region (Valdivia Villar, 1988; Elmoor-Loureiro, 2000; Zoppi de Roa, López, 2008; Kotov, Fuentes-Reinés, 2015). Although nowadays taxonomists consider this species as a complex of sibling taxa, *C. cf. cornuta* can be easily distinguished from other *Ceriodaphnia* species based on its prominent rostrum. Neotropical populations need to be revised in detail as part of a global revision of this group.

(2) *Ceriodaphnia dubia* Richard, 1894

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *C. dubia* is initially described from Indonesia, but then found in different regions of the world with tropical and temperate climate (see Chatterjee *et al.*, 2013). It has been reported from Andean lakes in nearby countries (Zoppi de Roa, López, 2008; De los Ríos Escalante, Kotov, 2015; Kotov, Fuentes-Reinés, 2015). This taxon needs a global revision, as it is apparently a group of closely related species (e.g. Sharma, 2014).

(3) *Daphnia* (*Daphnia*) *ambigua* Scourfield, 1947

Records. Andean Region: Lakes of Reserva Ecológica Cayambe-Coca (Páramo de Guamaní) (0°21'00"S 78°12'00"W) (Torres, Rylander, 2006). Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l. Province of Azuay (Alonso *et al.*, 2017). Lake Colta (1°45'00"S 78°44'33"W), 3.288 m a.s.l. (Villalobos, 1994).

Taxonomical remarks. The taxon was initially described from the United Kingdom, but the description was based on an exotic population occurring out of its main distribution range in the New World (see Benzie, 2005). This small-sized species of *Daphnia* is distributed throughout the Americas (Matsumura-Tundisi, 1984, Adamowicz *et al.*, 2004; Benzie, 2005) and invaded Europe. South American populations form a distinct clade (Hebert *et al.*, 2003). It could be confused with *D. parvula*, which has been also recorded from Ecuador.

(4) *Daphnia* (*Daphnia*) *laevis* Birge, 1879

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. This taxon was initially described from U.S.A. (see Benzie, 2005). It represents a group of cryptic species distributed in North and South America and Africa (Taylor *et al.*, 1998; Adamowicz *et al.*, 2004; Benzie, 2005).

(5) *Daphnia* (*Daphnia*) *longispina* (O.F. Müller, 1776)

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Labaj *et al.*, 2017; Van Colen *et al.*, 2017).

Taxonomical remarks. *D. longispina* was initially described from Denmark (see Kotov, 2015). This morphological confusing species complex has a wide geographical distribution in Africa, Europe and Asia (Benzie, 2005, Kotov *et al.*, 2013). Its taxonomy is confused even in Europe (Petrusek *et al.*, 2008; Kotov, 2015). As *D. longispina* s.str. does not occur in South America (Adamowicz *et al.*, 2004), this is most likely a misidentification.

(6) *Daphnia (Daphnia) cf. obtusa* Kurz, 1875

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *Daphnia obtusa* was initially described from Czech Republic (see Benzie, 2005). It is considered now a nearly cosmopolitan complex of species with numerous species (Benzie, 2005). Populations assigned to *D. obtusa* s.l. have been reported from high altitude regions of different tropical countries (Villalobos, 1994; Kotov, Taylor, 2010; Van Damme, Eggermont, 2011). Populations are known from the Neotropics (Villalobos, 1994). In South America this species needs to be revised, it is apparently represented by a series of taxa even on this continent (Kotov, Taylor, 2010).

(7) *Daphnia (Daphnia) parvula* (Fordyce, 1901)

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *D. parvula* has been described for North America, but has a wide geographical range in the western hemisphere, especially latitudinal (Benzie, 2005; Popova, Kotov, 2013), and widely distributed in Europe (Alonso, 1996; Flössner, 2000). This taxon could be confused with *D. ambigua*, which is also present in our checklist.

(8) *Daphnia (Daphnia) cf. pulex* Leydig, 1860

Records. Andean Region: Lake San Pablo (0°12'00"N 78°13'00"E), 2.660 m.a.s.l., Province of Imbabura (Gunkel, 2000; Labaj *et al.*, 2017). Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. It was initially described from unknown locality in Europe (Benzie, 2005). This taxon is a complex of species and a revision of the group is needed (Kotov, Taylor, 2010; Kotov, 2016), in particular for populations from South America (Crease *et al.*, 2012) and may be confused with *D. pulicaria*.

(9) *Daphnia (Daphnia) pulicaria* Forbes, 1893

Records. Andean Region: Lake San Pablo (0°12'00"N 78°13'00"E), 2.660 m a.s.l., Province of Imbabura (Villalobos, 1998; Casallas, Gunkel, 2001). Lake Coicochoa (0°18'00"N 78°21'00"W), 3.246 m a.s.l. (Villalobos, 1994). Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. This taxon was initially described from U.S.A. (see Benzie, 2005). According to Adamowicz *et al.* (2004) some records of *D. pulex* in South America may refer to *D. pulicaria*, and both taxa are easily confused also in Europe (see also in Crease *et al.* (2012)).

(10) *Simocephalus acutifrons* Johnson, 1953

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m.a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. This species was described from the Republic of South Africa (see Orlova-Bienkowskaja, 2001). Then it was considered as a junior synonym of *S. brehmi* Gauthier, 1939 (Orlova-Bienkowskaja, 2001; Kotov *et al.*, 2013). Both taxa were described from the African continent (Gauthier, 1939; Johnson, 1953) and populations of *S. cf. brehmi* have been reported from Brazil (Orlova-Bienkowskaja, 2001; Van Damme, Dumont, 2010). This is a very easily recognizable taxon, yet the status in South American populations needs revision (Van Damme, Dumont, 2010). The Ecuadorian record needs revision in comparison to other Neotropical populations of *S. cf. brehmi*.

(11) *Simocephalus exspinosus* (De Geer, 1778)

Records. Galápagos Islands: Lake El Junco (0°53'42"S 89°28'48"W), 675 m a.l.s., San Cristóbal Island (Obando, 2009).

Taxonomical remarks. The species was initially described from Europe, now it is considered as cosmopolitan taxon (Orlova-Bienkowskaja, 2001) which needs a global revision.

(12) *Simocephalus cf. vetulus* (O.F. Müller, 1776)

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m.a.s.l., Prov-

ince of Guayas (Briones, 2012). Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *S. vetulus* is a Palearctic species initially described from Denmark (see Orlova-Bienkowskaja, 2001; Huang *et al.*, 2014). Unfortunately, Orlova-Bienkowskaja (2001) did not analyze material from South America. According to Kotov *et al.* (2010), *S. vetulus* or a very closely related species is present in this continent. Neotropical populations of this taxon need to be checked morphologically and genetically, keeping in mind the existence of numerous *vetulus*-like species in other regions of the world (Huang *et al.*, 2014).

Family Moinidae Goulden, 1968

(1) *Moina micrura* Kurz, 1874

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. The species was initially described from Czech Republic (see Petrusek *et al.*, 2004). This taxon represents a cosmopolitan group which needs a detailed global revision (Elías-Gutiérrez *et al.*, 1999; Petrusek *et al.*, 2004; Bekker *et al.*, 2016).

Family Ilyocryptidae Smirnov, 1976 sensu Smirnov, 1992

(1) *Ilyocryptus spinifer* Herrick, 1882

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. This species was initially described from U.S.A. (see Kotov, Dumont, 2000). It is a common tropicopolitan taxon, usually found in different localities in South America (Kotov, Dumont, 2000).

(2) *Ilyocryptus spinosus* (Štifter, 1988)

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (recorded by Alonso *et al.*, 2017).

Taxonomical remarks. This species was initially described from Germany (see Kotov, Štifter, 2006). *I. spinosus* has been reported from the Northwest of U.S.A, Northeast of Canada and the North of Europe (Kotov, Štifter,

2006). Ecuadorian populations should be compared morphologically and genetically with Nearctic and Palearctic populations. The former could belong to a described taxon related to *I. spinosus* s.str.

Family Macrothricidae Norman and Brady, 1867

(1) *Macrothrix* cf. *hirsuticornis* (Norman and Brady, 1867)

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. This species was initially described from the United Kingdom (see Kotov, 2007). According to Smirnov (1992), only Palearctic populations could be regarded as *M. hirsuticornis* s.str., yet this taxon was recorded from North and South America as well. Kotov (2007) concluded that all populations from southern South America belong to *M. oviformis* Ekman, 1900. The record from Ecuador and others from the Neotropics must be revised because its identity with *M. oviformis* is not obvious.

(2) *Drepanothrix dentata* Eurén, 1861

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. This species was initially described from Sweden (Smirnov, 1992). It is distributed in the Palearctic region only (Alonso, 1996; Flössner, 2000). This taxon may also be present in the Nearctic region (Kotov *et al.*, 2013). *Drepanothrix dentata* was considered by Kotov (2016) as a complex of species which needs to be revised taxonomically. Only Iglesias *et al.* (2016) recorded this species in South America previously. There is a high chance that populations from South America belong to a new species if it is not a misidentification.

Family Bosminidae Baird, 1845 sensu Sars, 1865

(1) *Bosmina* (*Bosmina*) *longirostris* (O.F. Müller, 1776)

Records. Andean Region: Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. This taxon was initially described from Denmark; preliminary genetic data suggests that this is a really cosmopolitan taxon (Chatterjee *et al.*, 2013).

(2) *Bosmina (Liederobosmina) huaronensis* Delachaux, 1918

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Remarks. This species is initially described from Peruvian Andes; it is widely distributed in the Americas (De los Ríos Escalante, Kotov, 2015).

Family Chydoridae Dybowski and Grochowski, 1894

(1) *Acroperus tupinamba* Sinev & Elmoor-Loureiro 2010

Records. Coastal Region: Lake Tembladera (3°29'29"S 79°59'40"W), 17 m a.s.l., Province of El Oro, based on remains in lake sediments (López-Blanco, Sinev, 2016).

Remarks. *A. tupinamba* was described from Brazil; it has a wide distribution in Brazil and likely in other areas in the Neotropics (Sinev, Elmoor-Loureiro, 2010).

(2) *Alona cf. affinis* (Leydig, 1860)

Records. Andean Region: Lakes of Reserva Ecológica Cayambe-Coca (Páramo de Guamaní) (0°21'00"S 78°12'00"W) (Torres, Rylander 2006).

Taxonomical remarks. The species was initially described from Germany (see Sinev, 1997). Any records from the New World are unlikely belong to European *A. affinis* s.str. To date, several New World populations have been assigned to existing as well as new names (Sinev, 1997; Van Damme *et al.*, 2010). In Lake Titicaca (high altitudes of Peru), an *A. affinis*-like population has been given a name yet needs a formal redescription (see Van Damme *et al.*, 2010). The record from Ecuador needs a revision; it probably refers to *A. ossiani* (Sinev, 1998). In this checklist *A. affinis* is considered a dubious record.

(3) *Alona glabra* Sars, 1901

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–

4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. The taxon was initially described from Argentina; this is a widespread Neotropical species (Sinev, 2001). But it deserves close attention, because there are at least two different phylogroups in the region (Elías-Gutiérrez *et al.*, 2008). Sinev (2015a) recently revised this group (including Neotropical taxa) and suggested to move all *A. pulchella*-related species to the genus *Ovalona*, proposing a new combination *Ovalona glabra* (Sars, 1901).

(4) *Alona guttata* Sars, 1862

Records. Andean Region: Lakes of Reserva Ecológica Cayambe-Coca (Páramo de Guamaní) (0°21'00"S 78°12'00"W) (Torres, Rylander, 2006). Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Labaj *et al.*, 2017).

Taxonomical remarks. This taxon was initially described from Norway; it represents a cosmopolitan complex of species, widely distributed, which needs to be revised worldwide (Van Damme *et al.*, 2010). Neotropical populations show differences from North American species in this complex, *A. barbulata*, and from the European *A. guttata* (Sousa *et al.*, 2016).

(5) *Alona ossiani* (Sinev, 1998)

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. The taxon was described from Brazil (Sinev, 1998). *A. ossiani* is a Neotropical congener of the truly Palearctic *A. affinis* (Leydig, 1860) (Sinev, 1998; Van Damme *et al.*, 2010; Sinev, 2013). The previous record of *A. affinis* from Ecuador reported by Torres and Rylander (2006) most likely belongs to *A. ossiani* or to another related species, not to *A. affinis* s.str., and should be checked in detail.

(6) *Alpinalona cajasi* Alonso & Sinev, 2017

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso, Sinev, 2017).

Taxonomical remarks. Recorded initially by Alonso *et al.* (2017), as *Alona* sp. related to

Alona manueli Sinev et Zawisza, 2013 (*Alona* sp. gr. *manueli*) (Sinev, Zawisza, 2013). It was described the same year as a new species in a new genus *Alpinalona* Alonso et Sinev, 2017.

(7) *Alonella hampelae* Alonso & Kotov 2017

Records. Andean Region: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l. (Alonso, Kotov, 2017).

Taxonomical remarks. This taxon was recently described from Lakes of Cajas National Park (Ecuador) by Alonso and Kotov (2017). It belongs to the *A. excisa*-group. All records of *A. cf. excisa* of South America need revision because other taxa could be present there (Alonso, Kotov, 2017).

(8) *Alonella nana* Baird, 1843

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (in Briones (2012) as "*Alona nana*").

Taxonomical remarks. This taxon was initially described from the United Kingdom (see Smirnov, 1971). Briones (2012) recorded "*Alona nana* Baird 1843" which is an erroneous combination, the author apparently meant *Alonella nana*. However, *A. nana* does not occur in the Neotropics (Smirnov, 1996) and it is possible that the record of Briones (2012) has been confused with a small Neotropical taxon *A. dadayi*, which is recognizable by a long rostrum. This record in the lowlands of Ecuador may be invalid.

(9) *Camptocercus dadayi* (Stingelin, 1913)

Records. Andean Region: Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *C. dadayi* is a well-defined Neotropical taxon initially described from Colombia (see Sinev, 2015b).

(10) *Chydorus cf. sphaericus* (O.F. Müller, 1776)

Records. Andean Region: Lakes of Reserva Ecológica Cayambe-Coca (Páramo de Guamaní) (0°21'00"S 78°12'00"W) (Torres, Rylander, 2006). Coastal Region: Lake Tembladera (3°29'29"S 79°59'40"W), 17 m a.s.l., Province of El Oro (López-Blanco, Sinev, 2016). Andean Re-

gion: Lakes of Cajas National Park (2°50'00"S 79°10'00"W), 3.160–4.450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017; Labaj *et al.*, 2017).

Taxonomical remarks. The species was initially described from Denmark (Smirnov, 1971). This is a very difficult cosmopolitan species complex with many taxa, as revealed by genetic methods (Belyaeva, Taylor, 2009; Kotov *et al.*, 2016). The name "*C. sphaericus*" is used for nearly every *Chydorus* around the world, and it is not certain that all these records belong to the *C. sphaericus* group. Any records of this difficult genus should be accompanied by drawings or photos for confirmation.

(11) *Ephemeroporus acanthodes* Frey, 1982

Records. Andean Region: Lakes of Reserva Ecológica Cayambe-Coca (Páramo de Guamaní) (0°21'00"S 78°12'00"W) (Torres, Rylander, 2006, misspelled as "*E. acanthoides*").

Remarks. The species was initially described from souther U.S.A., in South America, it have been found in the Peruvian Andes (Valdivia Villar, 1988), a Venezuelan Andean lake (López, 1993) and more recently in different water bodies in Brazil (Serafim *et al.*, 2003, Van Damme, Dumont, 2010) and Paraguay (Debastiani-Júnior *et al.*, 2015). Several species of *Ephemeroporus* are present in the Neotropics, identifications should be checked thoroughly.

(12) *Euryalona occidentalis* Sars, 1901

Records. Galápagos Islands: Lake El Junco (0°53'42"S 89°28'48"W), 675 m a.s.l., San Cristóbal Island (Obando, 2009).

Taxonomical remarks. It is a clearly recognizable Neotropical taxon initially described from Brazil (Paggi, 1980).

(13) *Kurzia cf. latissima* (Kurz, 1875)

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. It was initially described from Czech Republic, its distribution is restricted to the Afrotropics and the Palearctic (Hudec, 2000). South American records have been shown to belong to *K. polyspina* (Hudec, 2000; Elmoor-Loureiro, 2002). It is most likely

that identification of Briones 2012 is erroneous, and the author dealt with *K. polyspina*.

(14) *Leydigopsis brevirostris* Brehm, 1938

Records. Coastal Region: Chongón Reservoir (02°13'09"S 80°07'32"W), 54 m a.s.l., Province of Guayas (Briones, 2012).

Taxonomical remarks. This is a well-defined Neotropical species initially described from Brazil (Sinev, 2004; Van Damme, Sinev, 2013), and other species in this genus can be expected in Ecuador.

(15) *Magnospina dentifera* (Sars, 1901)

Records. Coastal Region: Lake Tembladera (3°29'29"S 79°59'40"W), 17 m a.s.l., Province of El Oro, listed in the bottom sediments (López-Blanco, Sinev, 2016).

Taxonomical remarks. This taxon was initially described from Brazil (Sinev *et al.*, 2004). Recently, the *Alona dentifera* group was moved to the genus *Magnospina* Sousa, Elmoor-Loureiro *et al.*, 2016. However, it was separated from *Coronatella*, with unclear diagnosis, and the combination may change again. This Neotropical taxon ("*Alona*" *dentifera*) is well defined and occurs from U.S.A to Argentina (Sinev *et al.*, 2004; Sousa *et al.*, 2016).

(16) *Notoalona sculpta* (Sars, 1901)

Records. Coastal Region: Lake Tembladera (3°29'29"S 79°59'40"W), 17 m a.s.l., Province of El Oro, listed based on fragments found in lake sediments (López-Blanco, Sinev, 2016).

Taxonomical remarks. This is a clearly recognizable Neotropical taxon initially described from Brazil (Rajapaksa, Fernando, 1987; Van Damme *et al.*, 2013).

(17) *Oxyurella longicaudis* (Birge, 1910)

Records. Coastal Region: Lake Tembladera (3°29'29"S 79°59'40"W), 17 m a.s.l., Province of El Oro based on fragments found in lake sediments (López-Blanco, Sinev, 2016).

Remarks. *O. longicaudis* is a common Neotropical species initially described from unknown localities in South America (Elmoor-Loureiro, 1998; Van Damme, Dumont, 2010).

(18) *Paralona pigra* (Sars, 1862)

Records. Andean Region: Lakes of National Park Cajas (2°50'00"S 79°10'00"W), 3,160–

4,450 m a.s.l., Province of Azuay (Alonso *et al.*, 2017).

Taxonomical remarks. *P. pigra* was initially described from Norway; it is a cosmopolitan species (Dumont, Smirnov, 1996; Smirnov, 1996) and needs to be revised worldwide.

Discussion

The checklist of recorded species of Ecuadorian cladocerans contains 34 valid species. The number of cladoceran species of Ecuador recorded to date is significantly lower when compared to those in checklists for other countries in South America, such as Peru (88 taxa), Colombia (101 taxa), Brazil (116 taxa) and Venezuela (115 taxa) (Valdivia Villar, 1988; Elmoor-Loureiro, 2000; Zoppi de Roa, López, 2008; Kotov, Fuentes-Reinés, 2015). The comparably "low" biodiversity of the cladocerans in Ecuadorian water bodies is herein considered an artifact resulting from insufficient past efforts in sampling and identification. However, there is a benefit from the low number of studies on Cladocera in Ecuador: the taxonomical confusion is small and has not accumulated over decades (or even centuries). It is possible to start from a firm and modern taxonomical basis and conduct new studies in the region.

Several taxa in this checklist may belong to undescribed species, or to groups of which the taxonomy needs to be resolved in detail in the Neotropics or worldwide. The checklist illustrates that we should be careful to apply "Holarctic" names to Neotropical populations in Ecuador, including in high altitudes. However, we should be realistic that also "Löffler Islands" — tropical high altitude aquatic habitats harboring relict populations originating from northern latitudes (Van Damme, Eggermont, 2011) — could be found in Ecuador. In addition, the possibility of true human-mediated introductions should not be excluded. It is clear however, that the majority of records deserve a closer look and a morphological and genetic comparison to populations from the *terra typica*.

Taxonomical studies of freshwater Cladocera in Ecuador are just starting and most of the

efforts have been carried out in the limnetic zones of lakes and ponds in the Andean Region. Habitats important for cladoceran biodiversity, such as littoral zones, temporary pools and swamps, as well as several areas such as the low and medium altitudes of Ecuador, should be explored. The water bodies of the lowlands of the Coastal Region and Galápagos Islands have been scarcely studied and those of the Amazonian Region remain unexamined (Fig. 1). According to Green (1995), high altitude lakes have lower biodiversity than water bodies from lowlands (but see Van Damme, Eggermont, 2011), yet a high endemism can be expected in the Andes (Kotov *et al.*, 2010). Therefore, more extensive studies of cladocerans, including the unexplored lowlands, and further explorations in the mountains with unique taxa, are necessary to improve such a situation. In addition, it will be interesting to learn what the volcanic Galápagos really harbor of cladoceran faunas. Despite a relatively small country in comparison to its neighboring countries, Ecuador has also an enormous biodiversity as a result of the heterogeneity of existing environments and the notorious altitudinal gradients. The diversity in habitats is currently not reflected in the Cladocera.

Presently, any attempts to analyze the distributional patterns of Cladocera in Ecuador would be uninformative due to the scarcity of adequate information. The occurrence of Cladocera species that are common in the Neotropics and that are known for example in Colombia and Peru, as well as new species, can be expected, especially in the diverse family Chydoridae. The description of new chydorids from Ecuador (Alonso, Kotov, 2017; Alonso, Sinev, 2017) clearly point in that direction, yet new taxa in other families cannot be excluded.

Recommendations

The low number of Cladocera species reported from Ecuador could be attributed to the insufficient number of cladoceran surveys and studies. More taxonomical and biogeographical studies of Ecuadorian cladocerans are needed, especially in regions which are poorly studied,

such as the Amazonian Region, Coastal Region and Galápagos Islands (Fig. 1). These field surveys must encompass littoral zones of large lakes and ponds, particularly where macrophytes are present. Special attention should be paid to sampling in the bottom layers and in-shore areas and among macrophytes in order to survey the species that live in these zones. Also, it is necessary to sample a greater variety of aquatic habitats, such as artificial reservoirs, wetlands, swamps, “albarradas” (cisterns), inundated rice fields, temporary pools, moss, flooded forest, hyporheic/phreatic zones and phytotelmata communities. The efforts of local specialists should be combined with taxonomical training under the leadership of trained experts in order to provide knowledge transfer on taxonomy of cladocerans. The analysis of biodiversity of resting eggs from sediments using the Sars’ Method (Van Damme, Dumont, 2010) may be used complementary to obtain a better estimation of biodiversity of cladocerans, to describe different life stages (e.g. males) and provide material for molecular studies.

In concordance with recent advances in the field, these studies must be reinforced by genetics, in order to accurately explore the genetic isolation between populations from different regions of the world, and within the region.

Acknowledgements

Corrections and suggestions from anonymous referees greatly improved an early version of the manuscript. Special thanks to Dr F.D.R. Sousa for his valuable comments on some species and help with some bibliographical references and Dr A.Y. Sinev for some critical comments. Thanks for M. Belén López for help with the figure and L. Dávalos-Lind for revision of the English spelling and grammar in earlier draft.

Manuscript writing (remarks on taxonomic status of some tropical taxa revealed in Ecuador, general text editing) for ANN was funded by RFBR according to the research project 18-34-00389 мол_a. AAK was supported by Russian Government Program of Competitive Growth of Kazan Federal University.

References

- Adamowicz S.J., Hebert P.D.N., Marinone M.C. 2004. Species diversity and endemism in the *Daphnia* of Argentina: a genetic investigation // Zoological Journal of the Linnean Society. Vol.140. P.171–205.
- Alonso M. 1996. Crustacea, Branchiopoda. Fauna Iberica 7. Crustacea Branchiopoda. Madrid: Museo Nacional de Ciencias Naturales. Consejo Superior de Investigaciones Científicas. 486 p.
- Alonso M., Kotov A.A. 2017. A new species *Alonella* Sars, 1862 (Crustacea: Cladocera: Chydoridae) from Ecuadorian Andes // Zootaxa. Vol.4290. No.3. P.581–590.
- Alonso M., Sinev A.Yu. 2017. Relocation of *Alona manueli* Sinev & Zawisza, 2013 and a new closely related species from the Ecuadorian Andes to the new genus *Alpinalona* (Cladocera, Chydoridae, Aloninae) // Zootaxa. Vol.4350. No.3. P.563–573.
- Alonso M., Mosquera P., Hampel H., Vázquez R. 2017. Branquiópodos de las masas de agua lacustres del Parque Nacional Cajas (Andes Australes, Ecuador): inventario y notas autoecológicas // Maskana. Vol.8. No.1. P.71–74.
- Bekker E.I., Karabanov D.P., Galimov Y.R., Kotov A.A. 2016. DNA barcoding reveals high cryptic diversity in the North Eurasian *Moina* species (Crustacea: Cladocera) // PLoS ONE. Vol.11. No.8. e0161737.
- Belyaeva M., Taylor D.J. 2008. Cryptic species within the *Chydorus sphaericus* species complex (Crustacea: Cladocera) revealed by molecular markers and sexual stage morphology // Molecular Phylogenetics and Evolution. Vol.50. P.534–546.
- Benzie J.A.H. 2005. The genus *Daphnia* (including *Daphniopsis*) (Anomopoda: Daphniidae). Guides to the identification of the microinvertebrates of the continental waters of the world Vol.21. Leiden: Kenobi Productions, Ghent and Backhuys Publishers. 376 p.
- Brehm V. 1956. Cladocera aus Venezuela // Gessner F., Vareschi V. (Hrsg.). Ergebnisse der Deutschen Limnologischen. Venezuela Expedition 1952. S.217–232.
- Briones C. 2012. Estudio preliminar de la diversidad de Cladóceros en el Lago Chongón, Guayas (Bachelor's thesis). Guayaquil–Ecuador: Universidad de Guayaquil. 51 p.
- Casallas J., Gunkel G. 2001. Algunos aspectos limnológicos de un lago altoandino: El Lago San Pablo, Ecuador // Limnetica. Vol.20. No.2. P.215–232.
- Chatterjee T., Kotov A.A., Van Damme K., Chandrasekhar S.V.A., Padhye S. 2013. An annotated checklist of the Cladocera (Crustacea: Branchiopoda) from India // Zootaxa. Vol.3667. No.1. P.1–89.
- Colinvaux P., Steinitz-Kannan M. 1980. Species richness and area in Galapagos and Andean lakes: equilibrium phytoplankton communities and a paradox of the zooplankton // Kerfoot W.C. (ed.). Evolution and ecology of zooplankton communities. Hanover, New Hampshire: University Press of New England. P.697–712.
- De los Ríos Escalante P., Kotov A.A. 2015. A checklist of Branchiopoda (Anostraca and Cladocera) of Chilean continental waters // Zootaxa. Vol. 4027. No.3. P.366–388.
- Crease T.J., Omilian A.R., Costanzo K.S., Taylor D.J. 2012. Transcontinental phylogeography of the *Daphnia pulex* species complex // PLoS ONE. Vol.7. No.10. e46620.
- Debastiani-Júnior J.R., Elmoor-Loureiro L.M.A., Nogueira M.G. 2015. High taxonomic resolution as a determinant on finding new species and new records in the Río de La Plata basin: a case on Chydoridae (Crustacea: Branchiopoda: Cladocera) // Nauplius. Vol.23. No.1. P.21–30.
- Delachaux T. 1919. Cladóceres des Andes Péruviennes // Mémoires de la Société Neuchâteloise des Sciences Naturelles. Vol.43. P.18–38.
- Dumont H.J., Smirnov N.N. 1996. Reinstatement of the genus *Paralona* Šrámek-Hušek et al., 1962, and assessment of morphological variability in the *Paralona pigra*-group // Hydrobiologia. Vol.333. P.29–36.
- Elías-Gutiérrez M., Ciro-Pérez J., Suárez-Morales E., Silva-Briano M. 1999. The freshwater Cladocera (Orders Ctenopoda and Anomopoda) of Mexico, with comments on selected taxa // Crustaceana. Vol.72. No.2. P.171–186.
- Elías-Gutiérrez M., Martínez Jerónimo F., Ivanova N.V., Valdez Moreno M., Hebert P.D.N. 2008. DNA barcodes for Cladocera and Copepoda from Mexico and Guatemala, highlights and new discoveries // Zootaxa. Vol.1839. P.1–42.
- Elmoor-Loureiro L.M.A. 1998. Branchiopoda. Freshwater Cladocera // Young P.S. (ed.). Catalogue of Crustacea of Brazil. Rio de Janeiro: Museu Nacional. (Série Livros No. 6). P.15–41.
- Elmoor-Loureiro L.M.A. 2000. Brazilian cladoceran studies: where do we stand? // Nauplius. Vol.8. No.1. P.117–131.
- Elmoor-Loureiro L.M.A. 2002. Occurrence of *Kurzia polyspina* Hudec (Crustacea, Anomopoda, Chydoridae) in Brazil // Revista Brasileira de Zoologia. Vol.19. No.1. P.305–307.
- Elmoor-Loureiro L.M.A. 2018. Cladóceros do Brasil: Famílias Chydoridae e Eurycercidae. Available at: <https://cladocera.wordpress.com>
- Flössner D. 2000. Die Haplopoda und Cladocera (ohne Bosminidae) Mitteleuropas. Leiden: Backhuys. 428 S.
- Gauthier H. 1939. Contribution à l'étude de la Faune Dulçaquicole de la région du Tchad et particulièrement des branchiopodes et des ostracodes // Bulletin de l'Institut Français d'Afrique Noire. Vol.1. No.1. P.110–244.
- Green J. 1995. Altitudinal distribution of tropical planktonic Cladocera // Hydrobiologia. Vol.307. P.75–84.
- Gunkel G. 2000. Limnology of an equatorial high mountain lake in Ecuador, Lago San Pablo // Limnological Ecology and Management of Inland Waters. Vol.30. No.2. P.113–120.
- Hebert P.D.N., Witt J.D.S., Adamowicz S.J. 2003. Phylogeographical patterning in *Daphnia ambigua*: Region-

- al divergence and intercontinental cohesion // *Limnology and Oceanography*. Vol.48. No.1. P.261–268.
- Huang X., Shi X., Kotov A.A., Gu F. 2014. Confirmation through genetic analysis of the existence of many local phylogenetic clades of the genus *Simocephalus* (Crustacea, Cladocera) in China // *PLoS ONE*. Vol.9. No.11. e112808.
- Hudec I. 2000. Subgeneric differentiation within *Kurzia* (Crustacea: Anomopoda: Chydoridae) and a new species from Central America // *Hydrobiologia*. Vol.421. P.165–178.
- Iglesias C., Bonecker C., Brandão L., Crispim M.C., Eskinazi-Sant'Anna E.M., Gerhard M., Laço Portinho J., Maia-Barbosa P., Panarelli E., Santangelo J.M. 2016. Current knowledge of South American cladoceran diapause: A brief review // *International Review of Hydrobiology*. Vol.101. No. 3–4. P.91–104.
- Ihering H. 1895. Os Crustaceos phyllopoetes do Brasil // *Revista do Museu Paulista*. Vol.1. P.165–180.
- Jeong H., Kotov A.A., Lee W. 2014. Checklist of the freshwater Cladocera (Crustacea: Branchiopoda) of South Korea // *Proceedings of the Biological Society of Washington*. Vol.127. No.1. P.216–228.
- Ji G.H., Xiang X.F., Chen S.Z., Yu G.L., Kotov A.A., Dumont H.J. 2015. Annotated Checklist of Chinese Cladocera (Crustacea: Branchiopoda). Part II. Order Anomopoda (families Macrotrichidae, Euryceridae and Chydoridae) // *Zootaxa*. Vol.4044. No.2. P.241–269.
- Johnson D.S. 1953. On some Cladocera from South African muds // *Annals and Magazine of Natural History*. Ser.12. Vol.6. No.72. P.923–928.
- Korovchinsky N.M. 1992. Sididae & Holopediidae (Crustacea: Daphniiformes) // *Guides to the identification of the microinvertebrates of the continental waters of the world*. Vol.3. The Hague: SPB Academic Publishing. 82 p.
- Korovchinsky N.M. 2004. [Cladocerans of the order Ctenopoda of the world fauna (morphology, taxonomy, ecology, biogeography)]. Moscow: KMK Sci. Press. 410 p. [In Russian].
- Korovchinsky N.M. 2013. Cladocera (Crustacea: Branchiopoda) of South East Asia: history of exploration, taxon richness and notes on zoogeography // *Journal of Limnology*. Vol.72. No.52. P.109–124.
- Kotov A.A., 2007. Revision of the *hirsuticornis*-like species of *Macrothrix* Baird, 1843 (Cladocera: Anomopoda: Macrotrichidae) from Subantarctic and temperate regions of the southern hemisphere // *Journal of Natural History*. Vol.41. Nos.41–44. P.2569–2620.
- Kotov A.A. 2015. A critical review of the current taxonomy of the genus *Daphnia* O.F. Müller, 1785 (Anomopoda, Cladocera) // *Zootaxa*. Vol.3911. No.2. P.184–200.
- Kotov A.A. 2016. Faunistic complexes of the Cladocera (Crustacea, Branchiopoda) of Eastern Siberia and the Far East of Russia // *Biology Bulletin*. Vol.43. No.9. P.970–987.
- Kotov A.A., Dumont H.J. 2000. Analysis of the *Ilyocryptus spinifer*-species group (Anomopoda, Branchiopoda), with description of a new species // *Hydrobiologia*. Vol.428. No.1. P.85–113.
- Kotov A.A., Fuentes-Reinés J.M. 2015. An annotated checklist of the Cladocera (Crustacea: Branchiopoda) of Colombia // *Zootaxa*. Vol.4044. No.4. P.493–510.
- Kotov A.A., Štifter P. 2006. Cladocera: Family Ilyocryptidae (Branchiopoda: Cladocera: Anomopoda) // *Guides to the identification of the microinvertebrates of the Continental Waters of the world*. Vol.22. Leiden: Kenobi Productions, Ghent and Backhuys Publishers. 172 p.
- Kotov A.A., Taylor D.J. 2010. A new African lineage of the *Daphnia obtusa* group (Cladocera: Daphniidae) disrupts continental vicariance patterns // *Journal of Plankton Research*. Vol.32. No.6. P.937–949.
- Kotov A.A., Sinev A.Yu., Berrios V.L. 2010. The Cladocera (Crustacea: Branchiopoda) of six high altitude water bodies in the North Chilean Andes, with discussion of Andean endemism // *Zootaxa*. Vol.2430. No.1. P.1–66.
- Kotov A.A., Forró L., Korovchinsky N.M., Petrusek A. 2013. World checklist of freshwater Cladocera species. Available from: <https://fada.biodiversity.be/group/show/17>
- Kotov A.A., Karabanov D.P., Bekker E.I., Neterina T.V., Taylor D.J. 2016. Phylogeography of the *Chydorus sphaericus* group (Cladocera: Chydoridae) in the Northern Palearctic // *PLoS ONE*. Vol.11. No.12: e0168711. doi:10.1371/journal.pone.0168711
- Labaj A.L., Michelutti N., Smol J.P. 2017. Changes in cladoceran assemblages from tropical high mountain lakes during periods of recent climate change // *Journal of Plankton Research*. Vol.39. No.2. P.211–219.
- López C. 1993. Primer registro de *Ephemeroporus acanthodes* Frey 1982 (Crustacea: Cladocera) para Venezuela // *Acta Científica Venezolana*. Vol.44. P.312–314.
- López-Blanco C., Sinev A.Yu. 2016. Cladocera biodiversity in la Tembladera Lake (Ecuador): a palaeolimnological approach // *Crustaceana*. Vol.89. No.14. P.1611–1637.
- Matsumura-Tundisi T. 1984. Occurrence of species of the genus *Daphnia* in Brazil // *Hydrobiologia*. Vol.112. No.3. 161–165.
- Obando R. 2009. Evaluación de los parámetros físico-químicos y biológicos de la Laguna El Junco antes, durante y después de la erradicación de la tilapia (*Oreochromis niloticus*), Isla de San Cristóbal, Galápagos. Bachelor's thesis. Ecuador: Universidad del Azuay. 77 p.
- Orlova-Bienkowskaja M.Y. 2001. Daphniidae: genus *Simocephalus* // *Guides to the identification of the microinvertebrates of the continental waters of the World*. Vol.17. Leyden: Backhuys. 130 p.
- Paggi J.C. 1980. Aportes al Conocimiento de la fauna Argentina de Cladoceros. II. *Euryalona fasciculata* Daday, 1905 y *Euryalona occidentalis* Sars, 1901 // *Rev. Asoc. Cienc. Nat. Literal*. Vol.11. P.145–160.
- Petrusek A., Èrny M., Audenaert E. 2004. Large intercontinental differentiation of *Moina micrura* (Crustacea:

- Anomopoda): one less cosmopolitan cladoceran? // *Hydrobiologia*. Vol.526. P.73–81.
- Petrusek A., Hobýk A., Nilsson J.P., Skage M., Ěerný M., Brede N., Schwenk K. 2008. A taxonomic reappraisal of the European *Daphnia longispina* complex (Crustacea, Cladocera, Anomopoda) // *Zoologica Scripta*. Vol.37. P.507–519.
- Popova E.Y., Kotov A.A. 2013. Latitudinal patterns in the diversity of two subgenera of the genus *Daphnia* O.F. Müller (Crustacea: Cladocera: Daphniidae) // *Zootaxa*. Vol.3736. No.2. P.159–174.
- Quimi G.M. 2014. Composición y abundancia del zooplancton en la Represa San Vicente, Comuna Las Balsas, durante junio – noviembre del 2013, Santa Elena, Ecuador. Tesis de grado. Universidad Estatal Península de Santa Elena, Ecuador. 71 p. <http://repositorio.upse.edu.ec/handle/46000/1874>
- Rajapaksa R., Fernando C.H. 1987. Redescription and assignment of *Alona globulosa* Daday, 1898 to a new genus *Notoalona* and a description of *Notoalona freyi* sp. nov. // *Hydrobiologia*. Vol.144. P.131–153.
- Sars G.O. 1885. On some Australian Cladocera, raised from dried mud // *Forhandlinger i Videnskabs-Selskabet i Christiania*. Vol.8. P.1–46.
- Sars G.O. 1901. Contributions to the knowledge of the fresh-water Entomostraca of South America, as shown by artificial hatching from dried material. 1. Cladocera // *Archiv for Mathematik og Naturvidenskab*. Vol.23. No.3. P.1–102.
- Serafim J.M., Lansac-Tôha F.A., Paggi J.C., Velho L.F.M., Robertson B. 2003. Cladocera fauna composition in a river-lagoon system of the upper Paraná River floodplain, with a new record for Brazil // *Brazilian Journal of Biology*. Vol.63. No.2. P.349–356.
- Sharma P. 2014. Morphological and molecular identification of three *Ceriodaphnia* species (Cladocera: Daphniidae) from Australia // *Advances in Zoology*. Article ID 258134. 14 p.
- Sharma P., Kotov A.A. 2013. Molecular approach to identify sibling species of the *Ceriodaphnia cornuta* complex (Cladocera: Daphniidae) from Australia with notes on the continental endemism of this group // *Zootaxa*. Vol.3702. No.1. P.79–89.
- Sinev A.Yu. 1997. Review of the *affinis*-group of *Alona* Baird, 1843, with the description of a new species from Australia (Anomopoda Chydoridae) // *Arthropoda Selecta*. Vol.6. Nos.3–4. P.47–58.
- Sinev A.Yu. 1998. *Alona ossiani* sp.n., a new species of the *Alona affinis* complex from Brasil, deriving from the collection of G. O. Sars (Anomopoda Chydoridae) // *Arthropoda Selecta*. Vol.7. No.2. P.103–110.
- Sinev A.Yu. 2001. Redescription of *Alona glabra* Sars, 1901, a South American species of the *pulchella*-group (Branchiopoda: Anomopoda: Chydoridae) // *Arthropoda Selecta*. Vol.10. No.4. P.273–280.
- Sinev A.Yu. 2004. Redescription of two species of the genus *Leydigioopsis* Sars, 1901 (Branchiopoda, Anomopoda, Chydoridae) // *Invertebrate Zoology*. Vol.1. No.1. P.75–92.
- Sinev A.Yu. 2013. Cladocerans of *Alona affinis* group (Cladocera: Anomopoda: Chydoridae) from North America // *Zootaxa*. Vol. 3693. No.3. P.329–343.
- Sinev A.Yu. 2015a. Revision of the *pulchella*-group of *Alona* s. lato leads to its translocation to *Ovalona* Van Damme et Dumont, 2008 (Branchiopoda: Anomopoda: Chydoridae) // *Zootaxa*. Vol.4044. No.4. P.451–492.
- Sinev A.Yu. 2015b. Morphology and phylogenetic position of three species of genus *Camptocercus* Baird, 1843 (Cladocera: Anomopoda: Chydoridae) // *Zootaxa*. Vol.4040. No.2. P.169–186.
- Sinev A.Yu., Elmoor-Loureiro L.M.A. 2010. Three new species of chydorid cladocerans of subfamily Aloniinae (Branchiopoda: Anomopoda: Chydoridae) from Brazil // *Zootaxa*. Vol.2390. P.1–25.
- Sinev A.Yu., Zawisza E. 2013. Comments on cladocerans of crater lakes of the Nevado de Toluca Volcano (Central Mexico), with the description of a new species, *Alona manuei* sp. nov // *Zootaxa*. Vol.3647. No.2. P.390–400.
- Sinev A.Yu., Kotov A.A., Van Damme K. 2004. Morphology of a Neotropical cladoceran *Alona dentifera* (Sars, 1901), and its position within the Chydoridae Stebbing, 1902 (Branchiopoda: Anomopoda) // *Arthropoda Selecta*. Vol.13. No.3. P.99–107.
- Smirnov N.N. 1971. [Chydoridae of the world fauna] // *Fauna SSSR. Rakoobraznye*. Vol.1. Moscow: Nauka. 531 p. [In Russian].
- Smirnov N.N. 1992. The Macrothricidae of the world. Guides to the identification of the microinvertebrates of the Continental Waters of the world. Vol.1. The Hague: SPB Academic Publishing. 143p.
- Smirnov N.N. 1996. Cladocera: the Chydorinae and Sayciinae (Chydoridae) of the world // *Guides to the identification of the microinvertebrates of the Continental Waters of the world*. Vol.11. Amsterdam: SPB Academic Publishing. 197 p.
- Sousa F.D.R., Elmoor-Loureiro L.M.A., Santos S. 2016. Position of the *dentifera*-group in the *Coronatella*-branch and its relocation to a new genus: *Magnospina* gen. n. (Crustacea, Chydoridae, Aloniinae) // *ZooKeys*. Vol.586. P.95–119.
- Steere W. 1950. The phytogeography of Ecuador // *Studies in Ecuador Geography*. Vol.15. P.83–86.
- Stingelin T. 1913. Cladoceren aus den gebirgen von Kolumbien. *Memories de la Societe des Sciences Naturelles de Neuchatel*. Vol.5. P.600–638.
- Taylor D.J., Finston T.L., Hebert P.D.N. 1998. Biogeography of a widespread freshwater crustacean: pseudocongruence and cryptic endemism in the North American *Daphnia laevis* complex // *Evolution*. Vol.52. No.6. P.1648–1670.
- Torres L.E., Rylander K. 2006. Diversity and abundance of littoral cladocerans and copepods in nine Ecuadorian highland lakes // *Revista de biología tropical*. Vol.54. No.1. P.131–137.
- Valdivia Villar R.S. 1988. Lista de Cladóceros dulceacuícolas del Perú // *Amazoniana*. Vol.10. No.3. P.283–297.
- Van Colen W.R., Mosquera P., Vanderstukken M., Goirisk K., Carrasco M.C., Decaestecker E., Alonso M., León-Tamariz F., Muylaert K. 2017. Limnology and trophic

- status of glacial lakes in the tropical Andes (Cajas National Park, Ecuador) // *Freshwater Biology*. Vol.62. No.3. P.458–473.
- Van Damme K., Eggermont H. 2011. The Afromontane Cladocera (Crustacea: Branchiopoda) of the Rwenzori (Uganda–D. R. Congo): taxonomy, ecology and biogeography // *Hydrobiologia*. Vol.676. P.57–100.
- Van Damme K., Dumont H.J. 2010. Cladocera of the Lençóis Maranhenses (NE-Brazil): faunal composition and a reappraisal of Sars' Method // *Brazilian Journal of Biology*. Vol.70. No.3. P.755–779.
- Van Damme K., Sinev A.Yu. 2013. Tropical Amphipacific disjunctions in the Cladocera (Crustacea: Branchiopoda) // *Journal of Limnology*. Vol.72. No.s2. P.209–244.
- Van Damme K., Kotov A.A., Dumont H.J. 2010. A checklist of names in *Alona* Baird 1843 (Crustacea: Cladocera: Chydoridae) and their current status: an analysis of the taxonomy of a lump genus // *Zootaxa*. Vol.2330. P.1–63.
- Van Damme K., Maiphæ S., Sa-artrit P. 2013. Inland swamps in South East Asia harbour hidden cladoceran diversities: species richness and the description of new paludal Chydoridae (Crustacea: Branchiopoda: Cladocera) from Southern Thailand // *Journal of limnology*. Vol.72. No.s2. P.174–208.
- Villalobos L. 1994. Distribution of *Daphnia* in high mountain and temperate lakes of South America // *Internationale Vereinigung für theoretische und angewandte Limnologie*. Vol.25. No.4. P.2400–2404.
- Xiang X.F., Ji G.H., Chen S.Z., Yu G.L., Xu L., Han B.P., Kotov A.A., Dumont H.J. 2015. Annotated Checklist of Chinese Cladocera (Crustacea: Branchiopoda). Part I. Haplopoda, Ctenopoda, Onychopoda and Anomopoda (families Daphniidae, Moinidae, Bosminidae, Ilyocryptidae) // *Zootaxa*. Vol.3904. No.1. P.1–27.
- Zoppi de Roa E., López C. 2008. An updated checklist of inland Cladocera (Crustacea: Orders Ctenopoda and Anomopoda) from Venezuela // *Zootaxa*. Vol.1919. P.45–57.

Responsible editor K.G. Mikhailov