

Rising rural body-mass index is the main driver of the global obesity epidemic in adults

NCD Risk Factor Collaboration (NCD-RisC)*

Body-mass index (BMI) has increased steadily in most countries in parallel with a rise in the proportion of the population who live in cities^{1,2}. This has led to a widely reported view that urbanization is one of the most important drivers of the global rise in obesity^{3–6}. Here we use 2,009 population-based studies, with measurements of height and weight in more than 112 million adults, to report national, regional and global trends in mean BMI segregated by place of residence (a rural or urban area) from 1985 to 2017. We show that, contrary to the dominant paradigm, more than 55% of the global rise in mean BMI from 1985 to 2017—and more than 80% in some low- and middle-income regions—was due to increases in BMI in rural areas. This large contribution stems from the fact that, with the exception of women in sub-Saharan Africa, BMI is increasing at the same rate or faster in rural areas than in cities in low- and middle-income regions. These trends have in turn resulted in a closing—and in some countries reversal—of the gap in BMI between urban and rural areas in low- and middle-income countries, especially for women. In high-income and industrialized countries, we noted a persistently higher rural BMI, especially for women. There is an urgent need for an integrated approach to rural nutrition that enhances financial and physical access to healthy foods, to avoid replacing the rural undernutrition disadvantage in poor countries with a more general malnutrition disadvantage that entails excessive consumption of low-quality calories.

Being underweight or overweight can lead to adverse health outcomes. BMI—a measure of underweight and overweight—is rising in most countries². It is commonly stated that urbanization is one of the most important drivers of the worldwide rise in BMI because diet and lifestyle in cities lead to adiposity^{3–6}. However, such statements are typically based on cross-sectional comparisons in one or a small number of countries. Only a few studies have analysed how BMI is changing over time in rural and urban areas. The majority have been in one country,

over short durations, and/or in one sex and narrow age groups. The few studies that covered more than one country^{7–12} used at most a few dozen data sources and hence could not systematically estimate trends, and focused primarily on women of child-bearing age.

Data on how BMI in rural and urban populations is changing are needed to plan interventions that address underweight and overweight. Here, we report on mean BMI in rural and urban areas of 200 countries and territories from 1985 to 2017. We used 2,009 population-based studies of human anthropometry conducted in 190 countries (Extended Data Fig. 1), with measurements of height and weight in more than 112 million adults aged 18 years and older. We excluded data based on self-reported height and weight because they are subject to bias. For each sex, we used a Bayesian hierarchical model to estimate mean BMI by year, country and rural or urban place of residence. As described in the Methods, the estimated trends in population mean BMI represent a combination of (1) the change in the health of individuals due to change in their economic status and environment, and (2) the change in the composition of individuals that make up the population (and their economic status and environment).

From 1985 to 2017, the proportion of the world's population who lived in urban areas¹ increased from 41% to 55%. Over the same period, global age-standardized mean BMI increased from 22.6 kg m⁻² (95% credible interval 22.4–22.9) to 24.7 kg m⁻² (24.5–24.9) in women, and from 22.2 kg m⁻² (22.0–22.4) to 24.4 kg m⁻² (24.2–24.5) in men. The increase in mean BMI was 2.09 kg m⁻² (1.73–2.44) and 2.10 kg m⁻² (1.79–2.41) among rural women and men, respectively, compared to 1.35 kg m⁻² (1.05–1.65) and 1.59 kg m⁻² (1.33–1.84) in urban women and men. Nationally, change in mean BMI ranged from small decreases among women in 12 countries in Europe and Asia Pacific, to a rise of >5 kg m⁻² among women in Egypt and Honduras. The lowest observed sex-specific mean BMI over these 33 years was that of rural women in Bangladesh of 17.7 kg m⁻² (16.3–19.2) and rural men in




Fig. 1 | The difference between rural and urban age-standardized mean BMI in women. a, Difference in age-standardized mean BMI in 1985. **b,** Difference in age-standardized mean BMI in 2017. We did not estimate the difference between rural and urban areas for countries and territories in which the entire population live in areas classified as urban (Singapore,

Hong Kong, Bermuda and Nauru) or rural (Tokelau)—shown in grey. See Extended Data Fig. 2 for mean BMI at the national level and in rural and urban populations in 1985 and 2017. See Extended Data Fig. 6 for comparisons of the results between women and men.

*A list of authors and their affiliations appears in the online version of the paper.




Fig. 2 | The difference between rural and urban age-standardized mean BMI in men. a, Difference in age-standardized mean BMI in 1985. **b**, Difference in age-standardized mean BMI in 2017. We did not estimate the difference between rural and urban areas for countries and territories in which the entire population live in areas classified as urban (Singapore,

Hong Kong, Bermuda and Nauru) or rural (Tokelau)—shown in grey. See Extended Data Fig. 3 for mean BMI at the national level and in rural and urban populations in 1985 and 2017. See Extended Data Fig. 6 for comparison of results between women and men.

Ethiopia of 18.4 kg m^{-2} (17.0–19.9), both in 1985; the highest were 35.4 kg m^{-2} (33.7–37.1) for urban women and 34.6 kg m^{-2} (33.1–35.9) for rural men in American Samoa in 2017 (Extended Data Figs. 2, 3), representing a twofold difference.

In 1985, urban men and women in every country in east, south and southeast Asia, Oceania, Latin America and the Caribbean and a region that comprises central Asia, the Middle East and north Africa had a higher mean BMI than their rural peers (Figs. 1, 2). The urban–rural




Fig. 3 | Trends in age-standardized mean BMI by rural and urban place of residence. a, Trends are shown for women in each region. **b**, Trends are shown for men in each region. The lines show the posterior mean estimates and the shaded areas show the 95% credible intervals.

Table 1 | Contributors to the rise in mean BMI from 1985 to 2017

		Rural component		Urban component		Urbanization component	
		Absolute contribution (kg m ⁻²)	Percentage contribution (%)	Absolute contribution (kg m ⁻²)	Percentage contribution (%)	Absolute contribution (kg m ⁻²)	Percentage contribution (%)
Emerging economies							
Central Asia, Middle East and north Africa	Men	1.30 (0.96–1.64)	48 (41–54)	1.33 (1.02–1.65)	49 (44–54)	0.09 (0.06–0.12)	3 (2–5)
	Women	1.96 (1.57–2.33)	59 (54–64)	1.31 (0.95–1.69)	39 (34–44)	0.06 (0.03–0.09)	2 (1–3)
East and southeast Asia	Men	1.99 (1.62–2.37)	67 (63–71)	0.66 (0.53–0.80)	22 (20–24)	0.33 (0.26–0.39)	11 (9–14)
	Women	1.81 (1.36–2.26)	73 (67–80)	0.47 (0.32–0.64)	19 (16–22)	0.18 (0.10–0.26)	7 (4–11)
Latin America and the Caribbean	Men	0.86 (0.63–1.09)	31 (26–37)	1.73 (1.31–2.16)	63 (58–67)	0.17 (0.13–0.20)	6 (5–8)
	Women	1.29 (1.07–1.51)	38 (34–43)	2.01 (1.56–2.49)	60 (55–63)	0.06 (0.03–0.10)	2 (1–3)
Oceania	Men	2.24 (1.12–3.37)	90 (80–102)	0.24 (–0.03–0.51)	10 (–2–20)	0.00 (0.00–0.00)	0 (0–0)
	Women	2.41 (0.89–3.98)	81 (69–90)	0.53 (0.18–0.89)	19 (10–31)	0.00 (0.00–0.00)	0 (0–0)
South Asia	Men	1.99 (1.42–2.54)	86 (79–94)	0.20 (0.00–0.40)	8 (0–15)	0.12 (0.09–0.15)	5 (3–8)
	Women	2.18 (1.46–2.87)	80 (73–87)	0.36 (0.13–0.60)	13 (6–19)	0.19 (0.16–0.23)	7 (5–11)
Sub-Saharan Africa							
Sub-Saharan Africa	Men	1.14 (0.64–1.63)	64 (53–73)	0.39 (0.22–0.55)	22 (15–28)	0.23 (0.19–0.27)	14 (10–21)
	Women	1.37 (0.90–1.83)	57 (49–63)	0.58 (0.42–0.74)	24 (21–28)	0.45 (0.42–0.49)	19 (15–25)
High-income and other industrialized regions							
Central and eastern Europe	Men	0.59 (0.35–0.82)	35 (26–44)	1.10 (0.70–1.50)	65 (57–73)	0.00 (–0.01–0.01)	0 (–1–1)
	Women	0.14 (–0.19–0.45)	NR	0.13 (–0.45–0.69)	NR	–0.02 (–0.03–0.00)	NR
High-income Asia Pacific	Men	0.48 (0.37–0.59)	31 (25–37)	1.15 (0.84–1.46)	72 (68–75)	–0.04 (–0.08–0.00)	–2 (–6–0)
	Women	0.12 (–0.01–0.27)	NR	–0.02 (–0.38–0.36)	NR	–0.10 (–0.15 to –0.06)	NR
High-income western countries	Men	0.58 (0.47–0.69)	24 (22–27)	1.80 (1.53–2.07)	76 (74–78)	–0.01 (–0.02–0.00)	0 (–1–0)
	Women	0.39 (0.24–0.54)	21 (15–26)	1.44 (1.09–1.79)	79 (74–84)	0.00 (–0.02–0.01)	0 (–1–1)
World							
World	Men	1.24 (1.06–1.43)	57 (53–60)	0.65 (0.54–0.75)	30 (27–32)	0.30 (0.28–0.32)	14 (12–16)
	Women	1.22 (1.01–1.43)	60 (56–64)	0.56 (0.44–0.69)	28 (24–31)	0.25 (0.23–0.27)	13 (11–15)

Contributions of the rise in mean BMI in rural and urban populations and of urbanization to the rise in mean BMI from 1985 to 2017, by region. Urbanization is defined as an increase in the proportion of the population who live in urban areas. Percentage contributions were calculated as described in the Methods. The reported values are the means and 95% credible intervals. The three percentages sum to 100%. When one component causes an increase in BMI in a region and another does the opposite, the components can be negative or greater than 100%. Urban and rural mean BMI and the percentage of the population who live in urban areas in 1985 and 2017 for each region are provided in Extended Data Table 1. NR, percentage contribution was not reported, because the regional change in mean BMI (which appears in the denominator of the percentage contribution) was small (<0.5 kg m⁻²), leading to unstable estimates.

gap was as large as 3.25 kg m⁻² (2.57–3.96) in women and 3.05 kg m⁻² (2.44–3.68) in men in India. Over time, the BMI gap between rural and urban women shrank in all of these regions by at least 40%, as BMI rose faster in rural areas than in cities (Fig. 3). In 14 countries in these regions, including Armenia, Chile, Jamaica, Jordan, Malaysia, Taiwan and Turkey, the ordering of rural and urban female BMI reversed over time and rural women had higher BMI than their urban peers in 2017 (Fig. 1 and Extended Data Fig. 4).

The mean BMI of rural men also increased more than the mean BMI of urban men in south Asia and Oceania, shrinking the urban–rural BMI gap by more than half (Figs. 2, 3). In east and southeast Asia, Latin America and the Caribbean, and central Asia, the Middle East and north Africa, men in both rural and urban areas experienced a similar BMI increase and, therefore, the urban excess BMI did not change substantially over time.

In contrast to emerging economies, excess BMI among urban women became larger in sub-Saharan Africa (Fig. 3): from 2.59 kg m⁻² (2.21–2.98) in 1985 to 3.17 kg m⁻² (2.93–3.42) in 2017 (posterior probability of the observed increase being a true increase >0.999). This occurred because female BMI rose faster in cities than in rural areas in sub-Saharan Africa. This led to women in sub-Saharan African countries, especially those in west Africa, having the largest urban excess BMI of any country in 2017—for example, more than 3.35 kg m⁻² in Niger, Burkina Faso, Togo and Ghana (Fig. 1 and Extended Data Fig. 4). BMI increased at a similar rate in rural and urban men in sub-Saharan Africa, with the difference in 2017 (1.66 kg m⁻²; 1.37–1.94) being similar to 1985 (1.60 kg m⁻²; 1.13–2.07) (Fig. 2 and Extended Data Fig. 4).

BMI was previously lower in rural areas of low- and middle-income countries than in cities, both because rural residents had higher energy expenditure in their daily work—especially agriculture—and domestic activities, such as fuelwood and water collection^{13,14}, and because lower incomes in rural areas restricted food consumption¹⁵. In middle-income countries, agriculture is increasingly mechanized, cars are used for rural transport as income increases and road infrastructure improves, service and administrative jobs have become more common in rural areas, and some household tasks are no longer needed—for example, because homes have a water connection and use commercial fuels¹⁶. Furthermore, higher incomes as a result of economic growth allow more spending on food and hence higher caloric intake, disproportionately more in rural areas, where a substantial share of income was previously spent on food. Additionally, the consumption of processed carbohydrates may have increased disproportionately in rural areas where such foods have become more readily available through national and transnational companies^{9,17–21}. These changes, referred to as ‘urbanization of rural life’ by some researchers⁶, have contributed to a larger increase in rural BMI^{22,23}.

In contrast to other regions, urbanization in sub-Saharan Africa preceded significant economic growth²⁴. Subsistence farming remains common in Africa, and agriculture remains mostly manual; fuelwood—usually collected by women—is still the dominant fuel in rural Africa; and the use of cars for transportation is limited by poor infrastructure and poverty. In African cities, many people have service and office jobs, and mobility has become less energy-intensive owing to shorter travel distances and the use of cars and buses. Furthermore, urban markets where fresh produce is sold are increasingly replaced by commercially prepared and processed

foods from transnational and local industries and street vendors^{25–27}. These effects are exacerbated by limited time and space for cooking healthy meals and possibly perceptions of large weight as a sign of affluence^{28,29}.

In contrast to low- and middle-income regions, urban women in high-income western and Asia Pacific regions, and in central and eastern Europe, had slightly lower mean BMI than their rural peers in 2017 (Fig. 3). The rural excess BMI for women in these regions changed little from 1985 to 2017. Nationally, the excess BMI of rural women was largest in central and eastern European countries (for example, around 1 kg m⁻² or more in Belarus, Latvia and Czech Republic; Fig. 1 and Extended Data Fig. 4). Rural men in high-income western countries also had an excess BMI compared to urban men throughout the analysis period. The largest rural excess BMI for men in 2017 was seen in Sweden, Czech Republic, Ireland, Australia, Austria and the United States, which all had an excess BMI of 0.35 kg m⁻² or larger. In the high-income Asia Pacific region and in central and eastern Europe, rural and urban men had almost identical BMI throughout these three decades (Fig. 2 and Extended Data Fig. 4).

The lower urban BMI in high-income and industrialized countries reflects a growing rural economic and social disadvantage, including lower education and income, lower availability and higher price of healthy and fresh foods^{30,31}, less access to, and use of, public transport and walking than in cities^{32,33}, and limited availability of facilities for sports and recreational activity³⁴, which account for a significant share of overall physical activity in high-income and industrialized countries.

We also estimated how much of the overall rise in mean BMI since 1985 has been due to increases in BMI of rural and urban populations versus those attributable to urbanization (defined as an increase in the proportion of the population who live in urban areas), in each region and in the world as a whole. At the global level, 60% (56–64) of the rise in mean BMI from 1985 to 2017 in women and 57% (53–60) in men was due to increases in the BMI of rural populations; 28% (24–31) in women and 30% (27–32) in men due to the rise in BMI in urban populations; and 13% (11–15) and 14% (12–16) due to urbanization (Table 1). The contribution of the rise in rural BMI ranged from around 60% to 90% in the mostly rural regions of sub-Saharan Africa, east, south and south-east Asia and Oceania. The contribution of urbanization was small in all regions of the world, with maximum values of 19% (15–25) among women and 14% (10–21) among men in sub-Saharan Africa.

Our results show that, contrary to the prevailing view^{3–6}, BMI is rising at the same rate or faster in rural areas compared to cities, particularly in low- and middle-income countries except among women in sub-Saharan Africa. These trends have resulted in a rural–urban convergence in BMI in most low- and middle-income countries, especially for women. This convergence mirrors the experience of high-income and industrialized countries, where we found a persistently higher BMI in rural areas. The rising rural BMI is the largest contributor to the BMI rise in low- and middle-income regions and in the world as a whole over the last 33 years, which challenges the current paradigm of urban living and urbanization as the key driver of the global epidemic of obesity.

In poor societies, urban areas historically had lower levels of undernutrition^{35,36}, possibly because infrastructure such as roads and electricity facilitate food trade, transport and storage in cities, which can in turn reduce the impacts of agricultural shocks and seasonality. As economic growth and rural nutrition programmes reduce rural caloric deficiency, the rural undernutrition disadvantage may be replaced with a more general and complex malnutrition that entails excessive consumption of low-quality calories. To avoid such an unhealthy transition, the fragmented national and international responses to undernutrition and obesity should be integrated, and the narrow focus of international aid on undernutrition should be broadened, to enhance access to healthier foods in poor rural and urban communities.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, statements of data availability and associated accession codes are available at <https://doi.org/10.1038/s41586-019-1171-x>.

Received: 29 October 2018; Accepted: 30 March 2019;
Published online 8 May 2019.

- United Nations Department of Economic and Social Affairs, Population Division. *World Urbanization Prospects: the 2014 Revision*. <https://esa.un.org/unpd/wup/publications/files/wup2014-report.pdf> (United Nations, 2015).
- NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* **390**, 2627–2642 (2017).
- WHO. *Global Report on Urban Health: Equitable Healthier Cities for Sustainable Development*. Report No. 9241565276. <https://apps.who.int/iris/handle/10665/204715> (World Health Organization, 2016).
- Yusuf, S., Reddy, S., Öunpuu, S. & Anand, S. Global burden of cardiovascular diseases: part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* **104**, 2746–2753 (2001).
- Wagner, K. H. & Brath, H. A global view on the development of non communicable diseases. *Prev. Med.* **54**, S38–S41 (2012).
- Popkin, B. M. Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am. J. Clin. Nutr.* **84**, 289–298 (2006).
- Abubakari, A. R. et al. Prevalence and time trends in obesity among adult West African populations: a meta-analysis. *Obes. Rev.* **9**, 297–311 (2008).
- Filozof, C., Gonzalez, C., Sereday, M., Mazza, C. & Braguinsky, J. Obesity prevalence and trends in Latin-American countries. *Obes. Rev.* **2**, 99–106 (2001).
- Jaacks, L. M., Slining, M. M. & Popkin, B. M. Recent underweight and overweight trends by rural–urban residence among women in low- and middle-income countries. *J. Nutr.* **145**, 352–357 (2015).
- Mamun, A. A. & Finlay, J. E. Shifting of undernutrition to overnutrition and its determinants among women of reproductive ages in the 36 low to medium income countries. *Obes. Res. Clin. Pract.* **9**, 75–86 (2015).
- Neuman, M., Kawachi, I., Gortmaker, S. & Subramanian, S. V. Urban–rural differences in BMI in low- and middle-income countries: the role of socioeconomic status. *Am. J. Clin. Nutr.* **97**, 428–436 (2013).
- Popkin, B. M., Adair, L. S. & Ng, S. W. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr. Rev.* **70**, 3–21 (2012).
- Assah, F. K., Ekelund, U., Brage, S., Mbanya, J. C. & Wareham, N. J. Urbanization, physical activity, and metabolic health in sub-Saharan Africa. *Diabetes Care* **34**, 491–496 (2011).
- Levine, J. A. et al. The work burden of women. *Science* **294**, 812 (2001).
- Subramanian, S. & Deaton, A. The demand for food and calories. *J. Polit. Econ.* **104**, 133–162 (1996).
- Ng, S. W. & Popkin, B. M. Time use and physical activity: a shift away from movement across the globe. *Obes. Rev.* **13**, 659–680 (2012).
- Popkin, B. M. Nutrition, agriculture and the global food system in low and middle income countries. *Food Policy* **47**, 91–96 (2014).
- Reardon, T., Timmer, C. P. & Minten, B. Supermarket revolution in Asia and emerging development strategies to include small farmers. *Proc. Natl Acad. Sci. USA* **109**, 12332–12337 (2012).
- Mahajan, V. How Unilever reaches rural consumers in emerging markets. *Harv. Bus. Rev.* <https://hbr.org/2016/12/how-unilever-reaches-rural-consumers-in-emerging-markets> (2016).
- Jacobs, A. & Richtel, M. How big business got Brazil hooked on junk food. *The New York Times*. <https://www.nytimes.com/interactive/2017/09/16/health/brazil-obesity-nestle.html> (2017).
- Reardon, T., Timmer, C., Barrett, C. & Berdegue, J. The rise of supermarkets in Africa, Asia and Latin America. *Am. J. Agric. Econ.* **85**, 1140–1146 (2003).
- Ng, S. W., Norton, E. C. & Popkin, B. M. Why have physical activity levels declined among Chinese adults? Findings from the 1991–2006 China Health and Nutrition Surveys. *Soc. Sci. Med.* **68**, 1305–1314 (2009).
- Monda, K. L., Adair, L. S., Zhai, F. & Popkin, B. M. Longitudinal relationships between occupational and domestic physical activity patterns and body weight in China. *Eur. J. Clin. Nutr.* **62**, 1318–1325 (2008).
- Fay, M. & Opal, C. *Urbanization without Growth: A Not-So-Uncommon Phenomenon*. Policy Research Working Paper No. 241. <https://openknowledge.worldbank.org/bitstream/handle/10986/21373/wps2412.pdf?sequence=1&isAllowed=y> (World Bank, 2000).
- FAO. *Street Food in Urban Ghana*. <http://www.fao.org/3/a-i5804e.pdf> (Food and Agriculture Organization of the United Nations, 2016).
- Chilanga, E., Riley, L., Ngwira, J., Chalinda, C. & Masitala, L. *Food Insecurity in Informal Settlements in Lilongwe, Malawi*. <https://www.afsun.org/wp-content/uploads/2017/09/AFSUN25.pdf> (African Food Security Network, 2017).
- Tschirley, D., Reardon, T., Dolislager, M. & Snyder, J. The rise of a middle class in east and southern Africa: implications for food system transformation. *J. Int. Dev.* **27**, 628–646 (2015).
- Holdsworth, M., Gartner, A., Landais, E., Maire, B. & Delpuech, F. Perceptions of healthy and desirable body size in urban Senegalese women. *Int. J. Obes.* **28**, 1561–1568 (2004).
- Ettarh, R., Van de Vijver, S., Oti, S. & Kyobutungi, C. Overweight, obesity, and perception of body image among slum residents in Nairobi, Kenya, 2008–2009. *Prev. Chronic Dis.* **10**, 130198 (2013).
- Liese, A. D., Weis, K. E., Pluto, D., Smith, E. & Lawson, A. Food store types, availability, and cost of foods in a rural environment. *J. Am. Diet. Assoc.* **107**, 1916–1923 (2007).
- Lenardson, J. D., Hansen, A. Y. & Hartley, D. Rural and remote food environments and obesity. *Curr. Obes. Rep.* **4**, 46–53 (2015).

32. Ihara, M. et al. A cross-sectional study of the association between city scale and daily steps in Japan: Data from the National Health and Nutrition Survey Japan (NHNS-J) 2006–2010 (in Japanese). *Nippon Koshu Eisei Zasshi* **63**, 549–559 (2016).
33. Scheiner, J. A century of motorisation in urban and rural contexts: paths of motorisation in German cities. *Erdkunde* **66**, 313–328 (2012).
34. Seguin, R., Connor, L., Nelson, M., LaCroix, A. & Eldridge, G. Understanding barriers and facilitators to healthy eating and active living in rural communities. *J. Nutr. Metab.* **2014**, 146502 (2014).
35. Paciorek, C. J., Stevens, G. A., Finucane, M. M. & Ezzati, M. Children's height and weight in rural and urban populations in low-income and middle-income countries: a systematic analysis of population-representative data. *Lancet Glob. Health* **1**, e300–e309 (2013).
36. Krumdiek, C. L. The rural-to-urban malnutrition gradient. A key factor in the pathogenesis of urban slums. *J. Am. Med. Assoc.* **215**, 1652–1654 (1971).

Acknowledgements This study was funded by the Wellcome Trust. H.B. was supported by a Medical Research Council Doctoral Training Partnership Studentship, J.B. by a Royal Society Research Grant, and M.D.C. by an Academy of Medical Sciences Springboard Award. We thank L. Jaacks, B. Popkin, S. Sundberg and W. Willett for recommendations of relevant citations. The authors are responsible for the views expressed in this Letter and they do not necessarily represent the views, decisions, or policies of the institutions with which they are affiliated.

Reviewer information *Nature* thanks Miguel A. Martinez Beneito, Barry M. Popkin and the other anonymous reviewer(s) for their contribution to the peer review of this work.

Author contributions M.E. designed the study and oversaw research. H.B. led the data collection and statistical analysis, and prepared results. The other authors contributed to study design; collected, reanalysed, pooled and checked

data; analysed pooled data; and prepared results. M.E. and H.B. wrote the first draft of the manuscript with input from the other authors.

Competing interests M.E. reports a charitable grant from the AstraZeneca Young Health Programme, and personal fees from Prudential, Scor and Third Bridge, outside the submitted work. The other authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s41586-019-1171-x>.

Supplementary information is available for this paper at <https://doi.org/10.1038/s41586-019-1171-x>.

Reprints and permissions information is available at <http://www.nature.com/reprints>.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019

NCD Risk Factor Collaboration (NCD-RisC)

Honor Bixby¹, James Bentham², Bin Zhou¹, Mariachiara Di Cesare³, Christopher J. Paciorek⁴, James E. Bennett¹, Cristina Taddei¹, Gretchen A. Stevens⁵, Andrea Rodriguez-Martinez¹, Rodrigo M. Carrillo-Larco¹, Young-Ho Khang⁶, Maroje Sorić⁷, Edward W. Gregg⁸, J. Jaime Miranda⁹, Zulfiqar A. Bhutta^{10,11}, Stefan Savin⁵, Marisa K. Sophiea¹, Maria L. C. Iulilli¹, Bethlehem D. Solomon¹, Melanie J. Cowan⁵, Leanne M. Riley⁵, Goodarz Danaei¹², Pascal Bovet^{13,14}, Adela Chirita-Emandi¹⁵, Ian R. Hambleton¹⁶, Alison J. Hayes¹⁷, Nayu Ikeda¹⁸, Andre P. Kengne¹⁹, Avula Laxmaiah²⁰, Yanping Li²¹, Stephen T. McGarvey²¹, Aya Mostafa²², Martin Neovius²³, Gregor Starc²⁴, Ahmad A. Zainuddin²⁵, Leandra Abarca-Gómez²⁶, Ziad A. Abdeen²⁷, Shynar Abdrakhmanova²⁸, Suhaila Abdul Ghaffar²⁵, Zargar Abdul Hamid²⁹, Jamila Abubakar Garba³⁰, Niveen M. Abu-Rmeileh³¹, Benjamin Acosta-Cazares³², Robert J. Adams³³, Wichai Aekplakorn³⁴, Kaosar Afsana³⁵, Imelda A. Agdeppa³⁶, Carlos A. Aguilar-Salinas³⁷, Charles Agyemang³⁸, Mohamad Hasnan Ahmad²⁵, Noor Ani Ahmad²⁵, Naser Ahmad³⁹, Alireza Ahmadvand⁴⁰, Wolfgang Ahrens⁴¹, Kamel Ajlouni⁴², Fadia AlBuhairan⁴³, Shahla AlDhukair⁴⁴, Hazzaa M. Al-Hazzaa⁴⁵, Mohamed M. Ali⁵, Osman Ali⁴⁶, Ala'a Alkerwi⁴⁷, Amani Rashed Al-Othman⁴⁸, Rajaa Al-Raddadi⁴⁹, Mar Alvarez-Pedrerol⁵⁰, Eman Aly⁵¹, Deepak N. Amarpurkar^{52,576}, Philippe Amouey^{53,54}, Antoinette Amuzu⁵⁵, Lars Bo Andersen⁵⁶, Sigmund A. Anderssen⁵⁷, Lars H. Ångquist⁵⁸, Ranjit Mohan Anjana⁵⁹, Alireza Ansari-Moghaddam⁶⁰, Hajer Aounallah-Skhiri⁶¹, Joana Araújo⁶², Inger Ariansen⁶³, Tahir Aris²⁵, Raphael E. Arku⁶⁴, Nimmathota Arlappa²⁰, Krishna K. Aryal⁶⁵, Thor Aspelund⁶⁶, Felix K. Assah⁶⁷, Maria Cecília F. Assunção⁶⁸, May Soe Aung⁶⁹, Juha Auvinen⁷⁰, Mária Avdicová⁷¹, Ana Azevedo⁷², Fereidoun Azizi⁷³, Mehrdad Azmin⁴⁰, Bontha V. Babu⁷⁴, Azli Baharudin²⁵, Suhad Bahijiri⁴⁹, Jennifer L. Baker⁷⁵, Nagalla Balakrishna²⁰, Mohamed Bamoshmoosh⁷⁶, Maciej Banach⁷⁷, Piotr Bandosz⁷⁸, José R. Banegas⁷⁹, Carlo M. Barbagallo⁸⁰, Alberto Barceló⁸¹, Amina Barkat⁸², Aluisio J. D. Barros⁶⁸, Mauro V. G. Barros⁸³, Iqbal Bata⁸⁴, Anwar M. Batieha⁸⁵, Rosangela L. Batista⁸⁶, Zhamilya Battakova²⁸, Assembekov Batyrbek⁸⁷, Louise A. Baur¹⁷, Robert Beaglehole⁸⁸, Silvia Bel-Serrat⁸⁹, Habiba Ben Romdhane⁹⁰, Judith Benedics⁹¹, Mikhail Benet⁹², Salim Berkinbayev⁸⁷, Antonio Bernabe-Ortiz⁹, Gailute Bernotiene⁹³, Heloisa Bettio⁹⁴, Aroor Bhagyalaxmi⁹⁵, Sumit Bharadwaj⁹⁶, Santosh K. Bhargava⁹⁷, Hongsheng Bi⁹⁸, Yufang Bi⁹⁹, Anna Bieh⁶³, Elysée Claude Bika Lele¹⁰⁰, Mukharram Bikbov¹⁰¹, Bihungum Bista¹⁰², Dusko J. Bjelica¹⁰³, Peter Bjerregaard^{104,105}, Espen Bjertness¹⁰⁶, Marius B. Bjertness¹⁰⁶, Cecilia Björkelund¹⁰⁷, Anneke Blokstra¹⁰⁸, Simona Bo¹⁰⁹, Martin Bobak¹¹⁰, Lynne M. Boddy¹¹¹, Bernhard O. Boehm¹¹², Heiner Boeing¹¹³, Jose G. Boggia¹¹⁴, Carlos P. Boissonnet¹¹⁵, Marialaura Bonaccio¹¹⁶, Vanina Bongard¹¹⁷, Matthias Bopp¹¹⁸, Rossana Borchini¹¹⁹, Herman Borghs¹²⁰, Lien Braeckvelt¹²¹, Lutgart Braeckman¹²², Marjolijn C. E. Bragt¹²³, Imperia Brajkovich¹²⁴, Francesco Branca⁵, Juergen Breckenkamp¹²⁵, João Breda¹²⁶, Hermann Brenner¹²⁷, Lizzy M. Brewster³⁸, Garry R. Brian¹²⁸, Lacramioara Brinduse¹²⁹, Graziella Bruno¹⁰⁹, H. Bas Bueno-de-Mesquita¹⁰⁸, Anna Bugge¹³⁰, Marta Buoncristiani¹²⁶, Genc Burazeri¹³¹, Con Burns¹³², Antonio Cabrera de León¹³³, Joseph Cacciottolo¹³⁴, Hui Cai¹³⁵, Tilema Cama¹³⁶, Christine Cameron¹³⁷, José Camolas¹³⁸, Gamze Can¹³⁹, Günay Can¹⁴⁰, Ana Paula C. Cândido¹⁴¹, Felicia Cañete¹⁴², Mario V. Capanzana³⁶, Eduardo Capuano¹⁴³, Vincenzo Capuano¹⁴³, Viviane C. Cardoso⁹⁴, Axel C. Carlsson¹⁴⁴, Esteban Carmuega¹⁴⁵, Maria J. Carvalho¹⁴⁶, Felipe F. Casanueva¹⁴⁷, Juan-Pablo Casas¹¹⁰, Carmelo A. Caserta¹⁴⁸, Ertugrul Celikcan¹⁴⁹, Laura Censi¹⁵⁰, Juraci A. Cesar¹⁵¹, Snehalatha Chamukuttan¹⁵², Angelique W. Chan¹⁵³, Queenie Chan¹, Himanshu K. Chaturvedi¹⁵⁴, Nishi Chaturvedi¹¹⁰, Norsyamliana Che Abdul Rahim²⁵, Chien-Jen Chen¹⁵⁵, Fangfang Chen¹⁵⁶, Huashuai Chen¹⁵⁷, Shuohua Chen¹⁵⁸, Zhengming Chen¹⁵⁹, Ching-Yu Cheng¹⁵³, Yiling J. Cheng⁸, Angela Chetrit¹⁶⁰, Ekaterina Chikova-Iscener¹⁶¹, Arnau Chiolero¹⁶², Shu-Ti Chiou¹⁶³, María-Dolores Chirlaque¹⁶⁴, Belong Cho¹⁶⁵, Yumi Cho¹⁶⁶, Kaare Christensen¹⁰⁵, Diego G. Christofaro¹⁶⁷, Jerzy Chudek¹⁶⁸, Renata Cifkova^{169,170}, Michelle Cilia¹⁷¹, Eliza Cinteza¹⁷², Frank Claessens¹⁷³, Janine Clarke¹⁷⁴, Els Clays¹²², Hans Concini¹⁷⁵, Susana C. Confortin¹⁷⁶, Cyrus Cooper¹⁷⁷, Tara C. Coppinger¹³², Simona Costanzo¹¹⁶, Dominique Cotel¹⁷⁸, Chris Cowell¹⁷, Cora L. Craig¹³⁷, Amelia C. Crampin¹⁷⁹, Ana B. Crujeiras¹⁸⁰, Juan J. Cruz⁷⁹, Alexandra Cucu¹⁸¹, Liufu Cui¹⁵⁸, Jean Dallongeville¹⁷⁸, Albertino Damasceno¹⁸², Camilla T. Damsgaard⁵⁸, Rachel Dankner¹⁶⁰, Thomas M. Dantoft⁷⁵, Graziella D'Arrigo¹⁸³, Parasmani Dasgupta¹⁸⁴, Saeed Dastgiri¹⁸⁵, Luc Dauchet⁵⁴, Kairat Davletov⁸⁷, Guy De Backer¹²², Dirk De Bacquer¹²², Amalia De Curtis¹¹⁶, Giovanni de Gaetano¹¹⁶, Stefaan De Henauw¹²², Paula Duarte de Oliveira⁶⁸, Karin De Ridder¹⁸⁶, Susanne R. de Rooij¹⁸⁷, Delphine De Smedt¹²², Mohan Deepa⁵⁹, Alexander D. Deev¹⁸⁸, Abbas Dehghan¹⁸⁹, Hélène Delisle¹⁹⁰, Francis Delpuech¹⁹¹, Elaine Dennison¹⁷⁷, Valérie Deschamps¹⁹², Klodian Dhana¹⁸⁹, Meghnath Dhimal¹⁰², Augusto F. Di Castelnuovo¹⁹³, Juvenal Soares Dias-da-Costa¹⁹⁴, Alejandro Diaz¹⁹⁵, Zivka Dika⁷, Shirin Djalalinia¹⁹⁶, Ha T. P. Do¹⁹⁷, Annette J. Dobson¹⁹⁸, Maria Benedetta Donati¹¹⁶, Chiara Donfrancesco¹⁹⁹, Silvana P. Donoso²⁰⁰, Angela Döring²⁰¹, Maria Dorobantu¹⁷², Ahmad Reza Dorosty³⁹, Eleonora d'Orsi¹⁷⁶, Kouamelan Doua²⁰², Wojciech Drygas²⁰³, Jia Li Duan²⁰⁴, Charmaine A. Duante³⁶, Rosemary B. Duda²⁰⁵, Vesselka Duleva¹⁶¹, Virginija Dulskiene⁹³, Samuel C. Dumith¹⁵¹, Vilnis Dzerve²⁰⁶, Elzbieta Dziankowska-Zaborszczyk⁷⁷, Ricky Eddie²⁰⁷, Eruke E. Egbagbe²⁰⁸, Robert Eggertsen¹⁰⁷, Gabriele Eiben²⁰⁹, Ulf Ekelund⁵⁷, Jalila El Ati²¹⁰, Denise Eldemire-Shearer²¹¹, Marie Eliassen⁷⁵, Paul Elliott¹, Reina Engle-Stone²¹², Rajiv T. Erasmus²¹³, Cihangir Erem¹³⁹, Louise Eriksen¹⁰⁵, Johan G. Eriksson²¹⁴, Jorge Escobedo-de la Peña³², Alun Evans²¹⁵, David Faeh¹¹⁸, Caroline H. Fall¹⁷⁷, Victoria Farrugia Sant'Angelo¹⁷¹, Farshad Farzadfar³⁹, Mohammad R. Fattahi²¹⁶, Francisco J. Felix-Redondo²¹⁷, Trevor S. Ferguson²¹¹, Romulo A. Fernandes¹⁶⁷, Daniel Fernández-Bergés²¹⁸, Daniel Ferrante²¹⁹, Marika Ferrari²²⁰, Catterina Ferreccio²²¹, Eldridge Ferrer³⁶, Jean Ferrieres¹¹⁷, Anna Fijalkowska²²², Günther Fink^{223,224}, Sandra Fischer²²⁵, Eric Monterubio Flores²²⁶, Bernhard Föger¹⁷⁵, Leng Huat Foo²²⁷, Ann-Sofie Forslund²²⁸, Maria Forsner²²⁸, Heba M. Fouad⁵¹, Damian K. Francis²²⁹, Maria do Carmo Franco²³⁰, Oscar H. Franco¹⁸⁹, Guillermo Frontera²³¹, Flavio D. Fuchs²³², Sandra C. Fuchs²³³, Yuki Fujita²³⁴, Takuro Furusawa²³⁵, Zbigniew Gaciong²³⁶, Mihai Gafencu¹⁵, Daniela Galeone²³⁷, Fabio Galvano²³⁸, Jingli Gao¹⁵⁸, Manoli Garcia-de-la-Hera²³⁹, Dickman Gareta²⁴⁰, Sarah P. Garnett¹⁷, Jean-Michel Gaspoz²⁴¹, Magda Gasull²⁴², Louise Gaches²⁴³, Andrea Gazzinelli²⁴⁴, Harald Geiger¹⁷⁵, Johanna M. Geleijnse²⁴⁵, Ali Ghanbari³⁹, Erfan Ghasemi³⁹, Anoosheh Ghasemian⁴⁰, Oana-Florentina Gheorghe-Fronea¹⁷², Simona Giampaoli¹⁹⁹, Francesco Gianfagna^{246,193}, Tiffany K. Gil²⁴⁷, Jonathan Giovannelli⁵⁴, Glen Gironella³⁶, Aleksander Giwercman²⁴⁸, Justyna Godos²³⁸, Sibel Gogen¹⁴⁹, Rebecca A. Goldsmith²⁴⁹, David Goltzman²⁵⁰, Helen Gonçalves⁶⁸, Angel R. Gonzalez²⁵¹, David A. Gonzalez-Chica²⁴⁷, Marcela Gonzalez-Gross²⁵², Margot González-Leon³², Juan P. González-Rivas²⁵³, María-Elena González-Villalpando²⁵⁴, Frederic Gottrand⁵³, Antonio Pedro Graça²⁵⁵, Sidsel Graff-Iversen⁶³, Dušan Grafnetter²⁵⁶, Aneta Grajda²⁵⁷, Maria G. Grammatikopoulou²⁵⁸, Ronald D. Gregor⁸⁴, Tomasz Grodzicki²⁵⁹, Anders Grøntved¹⁰⁵, Giuseppe Grosso²³⁸, Gabriella Gruden¹⁰⁹, Dongfeng Gu²⁶⁰, Emanuela Gualdi-Russo²⁶¹, Elias F. Gudmundsson²⁶², Vilundur Gudnason⁶⁶, Ramiro Guerrero²⁶³, Idris Guessous²⁴¹, Andre L. Guimaraes²⁶⁴, Martin C. Gulliford²⁶⁵, Johanna Gunnlaugsdottir²⁶², Marc Guanter²⁶⁶, Xiuhua Guo²⁶⁷, Yin Guo²⁶⁷, Prakash C. Gupta²⁶⁸, Rajeev Gupta²⁶⁹, Oye Gureje²⁷⁰, Beata Gurzkowska²⁵⁷, Laura Gutierrez²⁷¹, Felix Gutzwiller¹¹⁸, Farzad Hadaegh⁷³, Charalambos A. Hadjigeorgiou²⁷², Rosa Haghshenas⁴⁰, Jytte Halkjær²⁷³, Rebecca Hardy¹¹⁰, Rachakulla Hari Kumar²⁰, Maria Hassapidou²⁷⁴, Jun Hata²⁷⁵, Teresa Haugsgjerd²⁷⁶, Jiang He²⁷⁷, Yuna He²⁷⁸, Regina Heidinger-Felso²⁷⁹, Mirjam Heinen⁸⁹, Tatjana Hejgaard²⁸⁰, Marleen Elisabeth Hendriks²⁸¹, Ana Henriques⁶², Leticia Hernandez Cadena²²⁶, Sauli Herralta²⁸², Victor M. Herrera²⁸³, Isabelle Herter-Aeberli²⁸⁴, Ramin Heshmat²⁸⁵, Allan G. Hill¹⁷⁷, Sai Yin Ho²⁸⁶, Suzanne C. Ho²⁸⁷, Michael Hobbs²⁸⁸, Albert Hofman¹⁸⁹, Wilma M. Hopman²⁸⁹, Andrea R. V. R. Horimoto²⁹⁰, Claudia M. Hormiga²⁹¹, Bernardo L. Horta⁶⁸, Leila Houti²⁹², Christina Howitt¹⁶, Thein Htay²⁹³, Aung Soe Htet²⁹⁴, Maung Maung Than Htike²⁹⁴, Yonghua Hu²⁹⁵, José María Huerta²⁹⁶, Ilpo Tapani Huhtaniemi¹, Constanta Huidumac Petrescu¹⁸¹, Martijn Huismans^{297,298}, Abdullatif Hussein³¹, Chinh Nguyen Huu¹⁹⁷, Inge Huybrechts²⁶⁶, Nahla Hwalla²⁹⁹, Jolanda Hyska¹³¹, Licia Iacoviello^{246,116}, Jesús M. Ibarluzea³⁰⁰, Mohsen M. Ibrahim³⁰¹, Norazizah Ibrahim Wong²⁵, M. Arfan Ikram¹⁸⁹, Vilma E. Irazola²⁷¹, Takafumi Ishida³⁰², Muhammad Islam¹⁰, Aziz al-Safi Ismail²²⁷, Vanja Ivkovic³⁰³, Masanori Iwasaki³⁰⁴, Tuia Jääskeläinen²¹⁴, Rod T. Jackson⁸⁸, Jeremy M. Jacobs³⁰⁵, Hashem Jaddou⁸⁵, Tazeen Jafar¹⁵³, Kenneth James²¹¹, Kazi M. Jamil⁴⁸, Konrad Jamrozik^{247,576}, Imre Janszky³⁰⁶, Edward Janus³⁰⁷, Juel Jarani³⁰⁸, Marjo-Riitta Jarvelin^{70,1}, Grazyna Jasienska²⁵⁹, Ana Jelakovic³⁰³, Bojan Jelakovic³⁰⁹, Garry Jennings³¹⁰, Seung-lyeal Jeong³¹¹, Chao Qiang Jiang³¹², Ramon O. Jimenez³¹³, Michel Joffres³¹⁴, Mattias Johansson²⁶⁶, Jari J. Jokelainen²⁸², Jost B. Jonas³¹⁵, Torben Jørgensen⁷⁵, Pradeep Joshi³¹⁶, Dragana P. Jovic³¹⁷, Jacek Józwiak³¹⁸, Anne Juolevi²¹⁴, Gregor Jurak²⁴, Vesna Jurek⁷⁹, Rudolf Kaaks¹²⁷, Anthony Kafatos³¹⁹, Eero O. Kajantie²¹⁴, Ofra Kalter-Leibovici¹⁶⁰, Nor Azmi Kamaruddin³²⁰, Yves Kameli¹⁹¹, Efthymios Kapantais³²¹, Khem B. Karki³²², Amir Kasaeian³⁹, Marzieh Katibeh³²³, Joanne Katz³²⁴, Peter T. Katzmarzyk³²⁵, Jussi Kauhanen³²⁶, Prabhdeep Kaur³²⁷, Maryam Kavousi¹⁸⁹, Gyullii Kazakbaeva¹⁰¹, Ulrich Keil³²⁸, Lital Keinan-Boker²⁴⁹, Sirkka Keinänen-Kiukaanniemi²⁸², Roya Kelishadi³²⁹, Cecily Kelleher⁸⁹, Han C. G. Kemper³³⁰, Alina Kerimkulova³³¹, Mathilde Kersting³³², Timothy Key¹⁵⁹, Yousef Saleh Khader⁸⁵, Davood Khalili⁷³

- Mohammad Khateeb⁴², Kay-Tee Khaw³³³, Bahareh Khairi⁷³, Alireza Khosravi³³⁴, Ilse M. S. L. Khouw¹²³, Stefan Kiechl³³⁵, Ursula Kiechl-Kohlendorfer³³⁵, Japhet Killewo³³⁶, Jeongseon Kim³³⁷, Yeon-Yong Kim³¹¹, Jeannette Klimont³³⁸, Jurate Klumbiene⁹³, Michael Knoflach³³⁵, Bhawesh Koirala³³⁹, Elin Kolle⁵⁷, Patrick Kolsteren¹²², Jürgen König³⁴⁰, Raija Korpelainen^{70,341}, Paul Korrovits³⁴², Magdalena Korzycka²²², Seppo Koskinen²¹⁴, Katsuyasu Kouda³⁴³, Viktoria A. Kovacs³⁴⁴, Sudhir Kowlessur³⁴⁵, Sławomir Koziol³⁴⁶, Wolfgang Kratzer³⁴⁷, Susi Kriemler¹¹⁸, Peter Lund Kristensen¹⁰⁵, Steinar Krokstad³⁰⁶, Daan Kromhout³⁴⁸, Herculina S. Kruger³⁴⁹, Ruzena Kubinova³⁵⁰, Renata Kuciene⁹³, Diana Kuh¹¹⁰, Urho M. Kujala³⁵¹, Enisa Kujundzic³⁵², Zbigniew Kulaga²⁵⁷, R. Krishna Kumar³⁵³, Marie Kunešová³⁵⁴, Pawel Kurjata²⁰³, Yadlapalli S. Kusuma³⁵⁵, Karl Kuulasmaa²¹⁴, Catherine Kyobutungi³⁵⁶, Quang Ngoc La³⁵⁷, Fatima Zahra Laamiri³⁵⁸, Tiina Laatikainen²¹⁴, Carl Lachat¹²², Youcef Laid³⁵⁹, Tai Hing Lam²⁸⁶, Maja Lang Morovic³⁶⁰, Vera Lanska²⁵⁶, Georg Lappas³⁶¹, Bagher Larijani³⁶², Tint Swe Latt³⁶³, Lars E. Laugsand³⁰⁶, Laura Lauria¹⁹⁹, Maria Lazo-Porras⁹, Khanh Le Nguyen Bao¹⁹⁷, Agnès Le Port³⁶⁴, Tuyen D. Le¹⁹⁷, Jeannette Lee³⁶⁵, Jeonghee Lee³³⁷, Paul H. Lee³⁶⁶, Terho Lehtimäki³⁶⁷, Daniel Lemogoum³⁶⁸, Naomi S. Levitt³⁶⁹, Christa L. Lilly³⁷⁰, Wei-Yen Lim³⁶⁵, M. Fernanda Lima-Costa³⁷¹, Hsien-Ho Lin³⁷², Xu Lin³⁷³, Lars Lind³⁷⁴, Allan Linneberg⁷⁵, Lauren Linnet¹⁰⁷, Mieczysław Litwin²⁵⁷, Jing Liu³⁷⁵, Helle-Mai Loit³⁷⁶, Luis Lopes¹⁴⁶, Tania Lopez³⁷⁷, Esther López-García⁷⁹, Roberto Lorbeer³⁷⁸, Paulo A. Lotufo⁹⁴, José Eugenio Lozano³⁷⁹, Dalia Luksiene⁹³, Annamari Lundqvist²¹⁴, Robert Lundqvist³⁸⁰, Nuno Lunet¹⁴⁶, Per Lytys³⁸¹, Guansheng Ma²⁹⁵, Jun Ma²⁹⁵, George L. L. Machado-Coelho³⁸², Aristides M. Machado-Rodrigues³⁸³, Suka Machi³⁸⁴, Stefania Maggi³⁸⁵, Dianna J. Magliano³⁸⁶, Emmanuella Magriplis³⁸⁷, Bernard Maire¹⁹¹, Marjeta Majer⁷, Marcia Makdisse³⁸⁸, Fatemeh Malekzadeh²¹⁶, Reza Malekzadeh²¹⁶, Rahul Malhotra¹⁵³, Sofia Malyutina³⁸⁹, Lynell V. Maniego³⁶, Yannis Manios³⁹⁰, Jim I. Mann³⁹¹, Enzo Manzato³⁹², Paula Margozzini²²¹, Anastasia Markaki³⁹³, Oonagh Markey³⁹⁴, Eliza Markidou Ioannidou³⁹⁵, Larissa Pruner Marques¹⁷⁶, Pedro Marques-Vidal³⁹⁶, Jaume Marrugat³⁹⁷, Rosemarie Martin³⁹⁸, Yves Martin-Prevel¹⁹¹, Reynaldo Martorell³⁹⁹, Eva Martos⁴⁰⁰, Stefano Marventano²³⁸, Shariq R. Masoodi⁴⁰¹, Ellisiv B. Mathiesen⁴⁰², Prashant Mathur⁴⁰³, Alicia Matijasevich⁹⁴, Tandi E. Matsha⁴⁰⁴, Artur Mazur⁴⁰⁵, Jean Claude N. Mbanya⁴⁶, Shelly R. McFarlane²¹¹, Martin McKee⁵⁵, Stela McLachlan⁴⁰⁶, Rachael M. McLean³⁹¹, Scott B. McLean¹⁷⁴, Breige A. McNulty⁸⁹, Safiah Md Yusof⁴⁰⁷, Sounnia Mediene-Benchechor²⁹², Jurate Medzioniene⁹³, Parinaz Mehdipour³⁹, Aline Meirhaeghe⁴⁰⁸, Jørgen Meisfjord⁶³, Christa Meisinger²⁰¹, Ana Maria B. Menezes⁶⁸, Geetha R. Menon⁷⁴, Gert B. M. Mensink⁴⁰⁹, Alibek Mereke⁸⁷, Indrapal I. Meshram²⁰, Andres Metspalu²²⁵, Haakon E. Meyer¹⁰⁶, Jie Mi¹⁵⁶, Kim F. Michaelsen⁵⁸, Nathalie Michels¹²², Kairit Mikkil²²⁵, Jody C. Miller³⁹¹, Cláudia S. Minderico⁴¹⁰, Juan Francisco Miquel²²¹, Daphne Mirkopoulou⁴¹¹, Erkin Mirrakhimov³³¹, Marjeta Misogoj-Durakovic⁷, Antonio Mistretta²³⁸, Veronica Mocanu⁴¹², Pietro A. Modesti⁴¹³, Sahar Saeedi Moghaddam³⁹, Bahram Mohajer³⁹, Mostafa K. Mohamed²², Kazem Mohammad³⁹, Noushin Mohammadi⁴¹⁴, Viswanathan Mohan⁵⁹, Salim Mohanna⁹, Muhammad Fadhli Mohd Yusoff²⁵, Farnam Mohebi³⁹, Marie Moitry^{415,416}, Drude Molbo⁵⁸, Line T. Møllehave⁷⁵, Niels C. Møller¹⁰⁵, Dénes Molnár²⁷⁹, Amirabbas Momenan⁷³, Charles K. Mondo⁴¹⁷, Eric A. Monterrubio⁴¹⁸, Kotsedi Daniel K. Monyeke⁴¹⁹, Jin Soo Moon⁴²⁰, Leila B. Moreira²³³, Alain Morejon⁴²¹, Luis A. Moreno⁴²², Karen Morgan⁴²³, Suzanne Morin²⁵⁰, Erik Lykke Mortensen⁵⁸, George Moschonis⁴²⁴, Malgorzata Mossakowska⁴²⁵, Jorge Mota¹⁴⁶, Anabela Mota-Pinto³⁸³, Mohammad Esmael Motlagh⁴²⁶, Jorge Motta⁴²⁷, Kelias P. Msyambo⁴²⁸, Thet Thet Mu⁴²⁹, Magdalena Muc³⁸³, Boban Mugoša³⁵², Maria Lorenza Muiesan⁴³⁰, Parvina Mukhtorova⁴³¹, Martina Müller-Nurasyid²⁰¹, Neil Murphy²⁶⁶, Jaakko Mursu³²⁶, Elaine M. Murtagh³⁹⁸, Sanja Music Milanovic^{360,7}, Vera Musil⁷, Iraj Nabipour⁴³², Shohreh Naderimaghani³⁹, Gabriele Nagel⁴³³, Balkish M. Naidu²⁵, Harunobu Nakamura⁴³⁴, Jana Námešná⁷¹, Ei Ei K. Nang³⁶⁵, Vinay B. Nangia⁴³⁵, Martin Nankap⁴³⁶, Sameer Narake²⁶⁸, Paola Nardone¹⁹⁹, Matthias Nauck³⁷⁸, Eva Maria Navarrete-Muñoz²³⁹, William A. Neal³⁷⁰, Keiu Nelis³⁷⁶, Liis Nelis³⁷⁶, Ilona Nenko²⁵⁹, Flavio Nervi²²¹, Chung T. Nguyen⁴³⁷, Nguyen D. Nguyen⁴³⁸, Quang Ngoc Nguyen⁴³⁹, Ramfis E. Nieto-Martínez⁴⁴⁰, Guang Ning⁹⁹, Toshiharu Ninomiya²⁷⁵, Sania Nishtar⁴⁴¹, Marianna Noale³⁸⁵, Oscar A. Noboa¹¹⁴, Teresa Norat¹, Sawada Norie⁴⁴², Davide Noto⁸⁰, Mohammad Al Nsour⁴⁴³, Eha Nurk³⁷⁶, Moffat Nyirenda⁵⁵, Galina Obreja⁴⁴⁴, Angélica M. Ochoa-Avilés²⁰⁰, Eiji Oda⁴⁴⁵, Kyungwon Oh¹⁶⁶, Kumiko Ohara²³⁴, Ryutaro Ohtsuka⁴⁴⁶, Örn Olafsson²⁶², Maria Teresa Anselmo Olinto⁴⁴⁷, Isabel O. Oliveira⁶⁸, Maciej Oltarzewski⁴⁴⁸, Mohd Azahadi Omar²⁵, Altan Onat¹⁴⁰, Terence W. O'Neill⁴⁴⁹, Sok King Ong⁴⁵⁰, Lariane M. Ono¹⁷⁶, Pedro Ordunez⁸¹, Dermot O'Reilly²¹⁵, Rui Ornelas⁴⁵¹, Ana P. Ortiz⁴⁵², Pedro J. Ortiz⁹, Merete Osler⁴⁵³, Clive Osmond⁴⁵⁴, Sergej M. Ostojic⁴⁵⁵, Afshin Ostovar³⁹, Johanna A. Otero²⁹¹, Kim Overvad³²³, Ellis Owusu-Dabo⁴⁵⁶, Fred Michel Paccaud⁴⁵⁷, Cristina Padez³⁸³, Ioannis Pagkalos²⁷⁴, Elena Pahomova²⁰⁶, Andrzej Pająk²⁵⁹, Domenico Palli⁴⁵⁸, Alberto Palloni⁴⁵⁹, Luigi Palmieri¹⁹⁹, Wen-Harn Pan¹⁵⁵, Songhomitra Panda-Jonas³¹⁵, Arvind Pandey¹⁵⁴, Francesco Panza⁴⁶⁰, Dimitrios Papandreou⁴⁶¹, Soon-Woo Park⁴⁶², Winsome R. Parnell³⁹¹, Mahboubeh Parsaeian³⁹, Ionela M. Pascanu⁴⁶³, Nikhil D. Patel⁴⁶⁴, Ivan Pecin^{309,303}, Mangesh S. Pednekar²⁶⁸, Nasheeta Peer⁴⁶⁵, Sergio Viana Peixoto³⁷¹, Markku Peltonen²¹⁴, Alexandre C. Pereira²⁹⁰, Cynthia M. Pérez⁴⁵², Napoleon Perez-Farinos⁴⁶⁶, Annette Peters²⁰¹, Astrid Petersmann³⁷⁸, Janina Petkeviciene⁹³, Ausra Petrauskienė⁹³, Niloofar Peykari¹⁹⁶, Son Thai Pham⁴⁶⁷, Daniela Pierannunzi¹⁹⁹, Iris Pigeot⁴⁶⁸, Hynek Pikhart¹¹⁰, Aida Pilav⁴⁶⁹, Lorenza Pilotto⁴⁷⁰, Francesco Pistelli⁴⁷¹, Freda Pitakaka⁴⁷², Aleksandra Piwonska²⁰³, Pedro Plans-Rubió⁴⁷³, Bee Koon Poh³²⁰, Hermann Pohlabein⁴⁶⁸, Raluca M. Pop⁴⁶³, Stevo R. Popovic¹⁰³, Michel Porta⁴⁷⁴, Marileen L. P. Portegies¹⁸⁹, Georg Posch¹⁷⁵, Dimitrios Poulimeneas²⁷⁴, Hamed Pouraram³⁹, Akram Pourshams⁴⁷⁵, Hossein Poustchi⁴⁷⁶, Rajendra Pradeepa⁵⁹, Alison J. Price⁵⁵, Jacqueline F. Price⁴⁰⁶, Jardena J. Puder³⁹⁶, Iveta Pudule⁴⁷⁷, Soile E. Puhakka^{341,70}, Maria Pui¹⁵⁵, Margus Punab³⁴², Radwan F. Qasrawi²⁷, Mostafa Qorbani⁴⁷⁸, Tran Quoc Bao⁴⁷⁹, Madhari S. Radhika²⁰, Ivana Radic⁴⁵⁵, Ricardas Radisauskas⁹³, Mahfuzar Rahman⁴⁸⁰, Mahmudur Rahman⁴⁸¹, Olli Raitakari⁴⁸², Manu Raj³⁵³, Hemalatha Rajkumar²⁰, Sherali Rakhmatulloev⁴³¹, Sudha Ramachandra Rao³²⁷, Ambady Ramachandran¹⁵², Jacqueline Ramke⁸⁸, Elisabete Ramos⁷², Rafael Ramos⁴⁸³, Lekhraj Rampal⁴⁸⁴, Sanjay Rampal⁴⁸⁵, Kodavanti Mallikharjuna Rao²⁰, Ramon A. Rascon-Pacheco³², Mette Rasmussen⁴⁸⁶, Josep Redon⁴⁸⁷, Paul Ferdinand M. Reganit⁴⁸⁸, Valéria Regecová⁴⁸⁹, Luis Revilla³⁷⁷, Lourdes Ribas-Barba⁴⁹⁰, Robespierre Ribeiro⁴⁹¹, Elio Riboli⁴, Fernando Rigo⁴⁹², Natascia Rinaldo²⁶¹, Tobias F. Rinke de Wit⁴⁹³, Ana Rito⁴⁹⁴, Raphael M. Ritti-Dias⁴⁹⁵, Juan A. Rivera²²⁶, Cynthia Robitaille⁴⁹⁶, Daniela Rodrigues³⁸³, Fernando Rodríguez-Artalejo⁷⁹, María del Cristo Rodríguez-Pérez⁴⁹⁷, Laura A. Rodríguez-Villamizar⁴⁹⁸, Rosalba Rojas-Martínez⁴¹⁸, Nipa Rojroongwasinkul³⁴, Dora Romaguera¹⁸⁰, Annika Rosengren^{499,107}, Ian Rouse⁵⁰⁰, Joel G. R. Roy¹⁷⁴, Adolfo Rubinstein²⁷¹, Frank J. Rühli¹¹⁸, Jean-Bernard Ruidavets¹¹⁷, Emma Ruiz Moreno⁵⁰¹, Blanca Sandra Ruiz-Betancourt³², Paola Russo⁵⁰², Petra Rust³⁴⁰, Marcin Rutkowski⁷⁸, Charumathi Sabanayagam⁵⁰³, Harshpal S. Sachdev⁵⁰⁴, Saeid Safiri⁵⁰⁵, Olfa Saidi⁹⁰, Benoit Salanave⁵⁹², Eduardo Salazar-Martínez²²⁶, Diego Salmerón²⁹⁶, Veikko Salomaa²¹⁴, Jukka T. Salonen⁵⁰⁶, Massimo Salvetti⁴³⁰, Jose Sánchez-Abanto⁵⁰⁷, Sandjaja⁵⁰⁸, Susana Sans⁵⁰⁹, Loreto Santa-Marina⁵¹⁰, Diana A. Santos⁵¹¹, Ina S. Santos⁶⁸, Osvaldo Santos⁵¹¹, Rute Santos¹⁴⁶, Sara Santos Sanz⁴⁶⁶, Jouko L. Saramies⁵¹², Luis B. Sardinha⁵¹¹, Nizal Sarrafzadegan⁵¹³, Kai-Uwe Saum¹²⁷, Savvas Savva²⁷², Mathilde Savy¹⁹¹, Marcia Scazufca⁵¹⁴, Angelika Schaffrath Rosario⁴⁰⁹, Herman Schargrodsky⁵¹⁵, Anja Schienkiewitz⁴⁰⁹, Karin Schindler⁵¹⁶, Sabine Schipf⁷⁸, Carsten O. Schmidt³⁷⁸, Ida Maria Schmidt⁵¹⁷, Ben Schöttker¹²⁷, Constance Schultz¹⁸⁷, Aletta E. Schutte^{349,19}, Sylvain Sebert⁷⁰, Aye Aye Sen²⁹⁴, Rusdidah Selamat²⁵, Vedrana Sember²⁴, Abhijit Sen³⁰⁶, Idowu O. Senbanjo⁵¹⁸, Sadaf G. Sepanlou³⁹⁹, Victor Sequeira¹⁴², Luis Serra-Majem⁵¹⁹, Jennifer Servais¹⁷⁴, Svetlana A. Shalnova¹⁸⁸, Sanjib K. Sharma³³⁹, Jonathan E. Shaw³⁸⁶, Lela Shengelia⁵²⁰, Kenji Shibuya³⁰², Hana Shimizu-Furusawa⁵²¹, Dong Wook Shin⁵²², Youchan Shin⁵⁰³, Alfonso Siano⁵⁰², Rosalynn Siantar⁵⁰³, Abba M. Sibai²⁹⁹, Antonio M. Silva⁸⁶, Diego Augusto Santos Silva¹⁷⁶, Mary Simon¹⁵², Judith Simons⁵²³, Leon A. Simons⁵²⁴, Khairil Si-Ramlee⁴⁵⁰, Agneta Sjöberg¹⁰⁷, Michael Sjöström²³, Jolanta Slowikowska-Hilczer⁷⁷, Przemysław Slusarczyk⁴²⁵, Liam Smeeth⁵⁵, Marieke B. Snijder³⁸, Hung-Kwan So²⁸⁶, Eugène Sobngwi⁶⁷, Stefan Söderberg²²⁸, Moesijanti Y. E. Soekatri⁵²⁵, Agustinus Soemantri⁵²⁶, Vincenzo Solfrizzi⁵²⁷, Emily Sonestedt²⁴⁸, Yi Song²⁹⁵, Thorkild I. A. Sørensen⁵⁸, Charles Sossa Jérôme⁵²⁸, Aïcha Soumaré⁵²⁹, Angela Spinelli¹⁹⁹, Igor Spiroski⁵³⁰, Jan A. Staessen⁵³¹, Hanspeter Stamm⁵³², Maria G. Stathopoulou⁵³³, Kaspar Staub¹¹⁸, Bill Stavreski³¹⁰, Jostein Steene-Johannessen⁵⁷, Peter Stehle⁵³⁴, Aryeh D. Stein³⁹⁹, George S. Stergiou⁵³⁵, Jochanan Stessman³⁰⁵, Doris Stöckl²⁰¹, Tanja Stocks²⁴⁸, Jakob Stokwiszewski⁵³⁶, Gareth Stratton⁵³⁷, Karen Stronks³⁸, Maria Wany Strufaldi²³⁰, Lela Sturua⁵²⁰, Ramón Suárez-Medina²⁵⁴, Chien-An Sun⁵³⁸, Johan Sundström³⁷⁴, Yn-Tz Sung²⁸⁷, Jordi Sunyer⁵⁰, Paibul Suriyawongpaisal³⁴, Boyd A. Swinburn⁸⁸, Rody G. Sy⁴⁸⁸, René Charles Sylva⁵³⁹, Lucjan Szponar⁴⁴⁸, E. Shyong Tai³⁶⁵, Mari-Liis Tammesoo²²⁵, Abdonas Tamosiunas⁹³, Eng Joo Tan¹⁷, Xun Tang²⁹⁵, Frank Tanser⁵⁴⁰, Yong Tao²⁹⁵, Mohammed Rasoul Tarawneh⁵⁴¹, Jakob Tarp⁵⁷, Carolina B. Tarqui-Mamani⁵⁰⁷, Radka Taxová Braunerová³⁵⁴, Anne Taylor²⁴⁷, Félicité Tchibindat⁴³⁶, William R. Tebar¹⁶⁷, Grethe Tell²⁷⁶, Tania Tello⁹, Holger Theobald¹⁴⁴, Xenophon Theodoridis²⁵⁸, Lutgarde Thijs⁵³¹, Betina H. Thuesen⁷⁵, Lubica Tichá⁵⁴², Erik J. Timmermans³³⁰, Anne Tjønneland²⁷³, Hanna K. Tolonen²¹⁴, Janne S. Tolstrup¹⁰⁵, Murat Topbas¹³⁹, Roman Topór-Madry²⁵⁹, María José Tormo⁵⁴³

- Michael J. Tornaritis²⁷², Maties Torrent⁵⁴⁴, Stefania Toselli⁵⁴⁵, Pierre Traissac¹⁹¹, Dimitrios Trichopoulos¹²⁵⁷⁶, Antonia Trichopoulou⁵⁴⁶, Oanh T. H. Trinh⁴³⁸, Atul Trivedi⁵⁴⁷, Yu-Hsiang Tsao³⁷², Lechaba Tshepo⁵⁴⁸, Maria Tsigga²⁷⁴, Shoichiro Tsugane⁴⁴², Marshall K. Tulloch-Reid²¹¹, Fikru Tullu⁵⁴⁹, Tomi-Pekka Tuomainen³²⁶, Jaakko Tuomilehto⁵⁵⁰, Maria L. Turley⁵⁵¹, Per Tynelius²³, Themistoklis Tzotzas³²¹, Christophe Tzourio⁵²⁹, Peter Ueda¹², Eunice E. Ugel⁵⁵², Flora A. M. Ukoi⁵⁵³, Hanno Ulmer³³⁵, Belgin Unal⁵⁵⁴, Hannu M. T. Uusitalo⁵⁵⁵, Justina Vaitkeviciute⁹³, Gonzalo Valdivia²²¹, Susana Vale⁵⁵⁶, Damaskini Valvi¹², Yvonne T. van der Schouw⁵⁵⁷, Koen Van Herck¹²², Hoang Van Minh³⁵⁷, Lenie van Rossem⁵⁵⁸, Natasja M. Van Schoor³³⁰, Irene G. M. van Valkengoed³⁸, Dirk Vanderschueren¹⁷³, Diego Vanuzzo⁴⁷⁰, Gregorio Varela-Moreiras⁵⁰¹, Patricia Varona-Pérez²⁵⁴, Lars Vatten³⁰⁶, Tomas Vega³⁷⁹, Toomas Veidebaum³⁷⁶, Gustavo Velasquez-Melendez²⁴⁴, Biruta Velika⁴⁷⁷, Giovanni Veronesi²⁴⁶, W. M. Monique Verschuren¹⁰⁸, Cesar G. Victora⁶⁸, Giovanni Viegi⁵⁵⁹, Lucie Viet¹⁰⁸, Paolo Vineis¹, Jesus Vioque⁵⁶⁰, Jyrki K. Virtanen³²⁶, Marjolein Visser²⁹⁸, Sophie Visvikis-Siest⁵³³, Koen Viswanathan⁵⁶¹, Tiina Vlasoff⁵⁶², Peter Vollenweider³⁹⁶, Henry Völzke³⁷⁸, Ari Voutilainen³²⁶, Sari Voutilainen³²⁶, Martine Vrijheid⁵⁰, Tanja G. M. Vrijkotte²⁹⁷, Alisha N. Wade⁵⁶³, Aline Wagner⁴¹⁶, Thomas Waldhör⁵¹⁶, Janette Walton¹³², Wan Mohamad Wan Bekakar²²⁷, Wan Nazaimoon Wan Mohamad⁵⁶⁴, Rildo S. Wanderley Jr⁸³, Ming-Dong Wang⁴⁹⁶, Qian Wang⁵⁶⁵, Xiangjun Wang⁵⁶⁶, Ya Xing Wang²⁶⁷, Ying-Wei Wang¹⁶³, S. Goya Wannamethee¹¹⁰, Nicholas Wareham³³³, Adelheid Weber⁹¹, Deepa Weerasekera⁵⁵¹, Daniel Weghuber⁵⁶⁷, Wenbin Wei²⁶⁷, Peter H. Whincup⁵⁶⁸, Kurt Widhalm⁵¹⁶, Indah S. Widyahening⁵⁶⁹, Andrzej Wiecek¹⁶⁸, Alet H. Wijga¹⁰⁸, Rainford J. Wilks²¹¹, Johann Willeit³³⁵, Peter Willeit³³⁵, Tom Wilsaard⁴⁰², Bogdan Wojtyniak⁵³⁶, Jyh Eiin Wong³²⁰, Tien Yin Wong¹⁵³, Roy A. Wong-McClure²⁶, Jean Woo²⁸⁷, Mark Woodward^{524,159}, Frederick C. Wu⁴⁴⁹, Jianfeng Wu⁹⁸, Shouling Wu¹⁵⁸, Haiquan Xu⁵⁷⁰, Liang Xu²⁶⁷, Uruwan Yamborisut³⁴, Weili Yan⁵⁷¹, Ling Yang¹⁵⁹, Xiaoguang Yang²⁷⁸, Yang Yang⁵⁶⁶, Nazan Yardim¹⁴⁹, Mehdi Yaseri⁷³, Xingwang Ye³⁷³, Panayiotis K. Yiallouros⁵⁷², Agneta Yngve³⁷⁴, Moein Yoosefi³⁹, Akihiro Yoshihara³⁰⁴, Qi Sheng You²⁶⁷, San-Lin You⁵³⁸, Novie O. Younger-Coleman²¹¹, Ahmad Faudzi Yusoff²⁵, Luciana Zaccagni²⁶¹, Vassilis Zafiropoulos³⁹³, Farhad Zamani⁵⁷³, Sabina Zambon³⁹², Antonis Zampelas³⁸⁷, Hana Zamrazilova³⁵⁴, Maria Elisa Zapata¹⁴⁵, Ko Ko Zaw³⁶³, Tomasz Zdrojewski⁷⁸, Tajana Zeljkovic Vrkic³⁰³, Yi Zeng^{157,295}, Dong Zhao³⁷⁵, Wenhua Zhao²⁷⁸, Wei Zheng¹³⁵, Yingfeng Zheng⁵⁰³, Bekbolat Zholdin⁵⁷⁴, Maigeng Zhou²⁷⁸, Dan Zhu⁵⁷⁵, Baurzhan Zhussupov⁸⁷, Esther Zimmermann⁷⁵, Julio Zúñiga Cisneros⁴²⁷ & Majid Ezzati^{1*}
- ¹Imperial College London, London, UK. ²University of Kent, Canterbury, UK. ³Middlesex University, London, UK. ⁴University of California Berkeley, Berkeley, CA, USA. ⁵World Health Organization, Geneva, Switzerland. ⁶Seoul National University, Seoul, South Korea. ⁷University of Zagreb, Zagreb, Croatia. ⁸US Centers for Disease Control and Prevention, Atlanta, GA, USA. ⁹Universidad Peruana Cayetano Heredia, Lima, Peru. ¹⁰Agá Khan University, Karachi, Pakistan. ¹¹The Hospital for Sick Children, Toronto, Ontario, Canada. ¹²Harvard T. H. Chan School of Public Health, Boston, MA, USA. ¹³Ministry of Health, Victoria, Seychelles. ¹⁴University of Lausanne, Lausanne, Switzerland. ¹⁵Victor Babeş University of Medicine and Pharmacy Timisoara, Timisoara, Romania. ¹⁶The University of the West Indies, Cave Hill, Barbados. ¹⁷University of Sydney, Sydney, New South Wales, Australia. ¹⁸National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo, Japan. ¹⁹South African Medical Research Council, Cape Town, South Africa. ²⁰ICMR-National Institute of Nutrition, Hyderabad, India. ²¹Brown University, Providence, RI, USA. ²²Ain Shams University, Cairo, Egypt. ²³Karolinska Institutet, Stockholm, Sweden. ²⁴University of Ljubljana, Ljubljana, Slovenia. ²⁵Ministry of Health Malaysia, Kuala Lumpur, Malaysia. ²⁶Caja Costarricense de Seguro Social, San José, Costa Rica. ²⁷Al-Quds University, East Jerusalem, Palestine. ²⁸National Center of Public Healthcare, Nur-Sultan, Kazakhstan. ²⁹Center for Diabetes and Endocrine Care, Srinagar, India. ³⁰Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria. ³¹Birzeit University, Birzeit, Palestine. ³²Instituto Mexicano del Seguro Social, Mexico City, Mexico. ³³Flinders University, Adelaide, South Australia, Australia. ³⁴Mahidol University, Nakhon Pathom, Thailand. ³⁵BRAC University, Dhaka, Bangladesh. ³⁶Food and Nutrition Research Institute, Taguig, The Philippines. ³⁷Instituto Nacional de Ciencias Médicas y Nutrición, Mexico City, Mexico. ³⁸University of Amsterdam, Amsterdam, The Netherlands. ³⁹Tehran University of Medical Sciences, Tehran, Iran. ⁴⁰Non-Communicable Diseases Research Center, Tehran, Iran. ⁴¹University of Bremen, Bremen, Germany. ⁴²The National Center for Diabetes, Endocrinology and Genetics, Amman, Jordan. ⁴³Aldara Hospital and Medical Center, Riyadh, Saudi Arabia. ⁴⁴King Abdullah International Medical Research Center, Riyadh, Saudi Arabia. ⁴⁵King Saud University, Riyadh, Saudi Arabia. ⁴⁶Universiti Malaysia Sabah, Kota Kinabalu, Malaysia. ⁴⁷Luxembourg Institute of Health, Strassen, Luxembourg. ⁴⁸Kuwait Institute for Scientific Research, Safat, Kuwait. ⁴⁹King Abdulaziz University, Jeddah, Saudi Arabia. ⁵⁰ISGlobal Centre for Research in Environmental Epidemiology, Barcelona, Spain. ⁵¹World Health Organization Regional Office for the Eastern Mediterranean, Cairo, Egypt. ⁵²Bombay Hospital and Medical Research Centre, Mumbai, India. ⁵³University of Lille, Lille, France. ⁵⁴Lille University Hospital, Lille, France. ⁵⁵London School of Hygiene & Tropical Medicine, London, UK. ⁵⁶Western Norway University of Applied Sciences, Sogndal, Norway. ⁵⁷Norwegian School of Sport Sciences, Oslo, Norway. ⁵⁸University of Copenhagen, Copenhagen, Denmark. ⁵⁹Madras Diabetes Research Foundation, Chennai, India. ⁶⁰Zahedan University of Medical Sciences, Zahedan, Iran. ⁶¹National Institute of Public Health, Tunis, Tunisia. ⁶²Institute of Public Health of the University of Porto, Porto, Portugal. ⁶³Norwegian Institute of Public Health, Oslo, Norway. ⁶⁴University of Massachusetts, Amherst, MA, USA. ⁶⁵Abt Associates, Kathmandu, Nepal. ⁶⁶University of Iceland, Reykjavik, Iceland. ⁶⁷University of Yaoundé 1, Yaoundé, Cameroon. ⁶⁸Federal University of Pelotas, Pelotas, Brazil. ⁶⁹University of Medicine 1, Yangon, Myanmar. ⁷⁰University of Oulu, Oulu, Finland. ⁷¹Regional Authority of Public Health, Banská Bystrica, Slovakia. ⁷²University of Porto Medical School, Porto, Portugal. ⁷³Shahid Beheshti University of Medical Sciences, Tehran, Iran. ⁷⁴Indian Council of Medical Research, New Delhi, India. ⁷⁵Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark. ⁷⁶University of Science and Technology, Sana'a, Yemen. ⁷⁷Medical University of Lodz, Lodz, Poland. ⁷⁸Medical University of Gdansk, Gdansk, Poland. ⁷⁹Universidad Autónoma de Madrid, Madrid, Spain. ⁸⁰University of Palermo, Palermo, Italy. ⁸¹Pan American Health Organization, Washington, DC, USA. ⁸²Mohammed V University de Rabat, Rabat, Morocco. ⁸³University of Pernambuco, Recife, Brazil. ⁸⁴Dalhousie University, Halifax, Nova Scotia, Canada. ⁸⁵Jordan University of Science and Technology, Irbid, Jordan. ⁸⁶Federal University of Maranhão, São Luís, Brazil. ⁸⁷Kazakh National Medical University, Almaty, Kazakhstan. ⁸⁸University of Auckland, Auckland, New Zealand. ⁸⁹University College Dublin, Dublin, Ireland. ⁹⁰University Tunis El Manar, Tunis, Tunisia. ⁹¹Federal Ministry of Labour, Social Affairs, Health and Consumer Protection, Vienna, Austria. ⁹²Cafam University Foundation, Bogota, Colombia. ⁹³Lithuanian University of Health Sciences, Kaunas, Lithuania. ⁹⁴University of São Paulo, São Paulo, Brazil. ⁹⁵B. J. Medical College, Ahmedabad, India. ⁹⁶Chirayu Medical College, New Delhi, India. ⁹⁷Sunder Lal Jain Hospital, Delhi, India. ⁹⁸Shandong University of Traditional Chinese Medicine, Shandong, China. ⁹⁹Shanghai Jiao-Tong University School of Medicine, Shanghai, China. ¹⁰⁰Institute of Medical Research and Medicinal Plant Studies, Yaoundé, Cameroon. ¹⁰¹Ufa Eye Research Institute, Ufa, Russia. ¹⁰²Nepal Health Research Council, Kathmandu, Nepal. ¹⁰³University of Montenegro, Niksic, Montenegro. ¹⁰⁴University of Greenland, Nuuk, Greenland. ¹⁰⁵University of Southern Denmark, Odense, Denmark. ¹⁰⁶University of Oslo, Oslo, Norway. ¹⁰⁷University of Gothenburg, Gothenburg, Sweden. ¹⁰⁸National Institute for Public Health and the Environment, Bilthoven, The Netherlands. ¹⁰⁹University of Turin, Turin, Italy. ¹¹⁰University College London, London, UK. ¹¹¹Liverpool John Moores University, Liverpool, UK. ¹¹²Nanyang Technological University, Singapore, Singapore. ¹¹³German Institute of Human Nutrition, Potsdam, Germany. ¹¹⁴Universidad de la República, Montevideo, Uruguay. ¹¹⁵CEMIC, Buenos Aires, Argentina. ¹¹⁶IRCCS Neuromed, Pozzilli, Italy. ¹¹⁷Toulouse University School of Medicine, Toulouse, France. ¹¹⁸University of Zurich, Zurich, Switzerland. ¹¹⁹University Hospital of Varese, Varese, Italy. ¹²⁰University Hospital KU Leuven, Leuven, Belgium. ¹²¹Flemish Agency for Care and Health, Brussels, Belgium. ¹²²Ghent University, Ghent, Belgium. ¹²³FrieslandCampina, Amersfoort, The Netherlands. ¹²⁴Universidad Central de Venezuela, Caracas, Venezuela. ¹²⁵Bielefeld University, Bielefeld, Germany. ¹²⁶World Health Organization Regional Office for Europe, Copenhagen, Denmark. ¹²⁷German Cancer Research Center, Heidelberg, Germany. ¹²⁸The Fred Hollows Foundation, Auckland, New Zealand. ¹²⁹University of Medicine and Pharmacy Bucharest, Bucharest, Romania. ¹³⁰University College Copenhagen, Copenhagen, Denmark. ¹³¹Institute of Public Health, Tirana, Albania. ¹³²Cork Institute of Technology, Cork, Ireland. ¹³³Universidad de La Laguna, Tenerife, Spain. ¹³⁴University of Malta, Pietà, Malta. ¹³⁵Vanderbilt University, Nashville, TN, USA. ¹³⁶Ministry of Health, Tongatapu, Tonga. ¹³⁷Canadian Fitness and Lifestyle Research Institute, Ottawa, Ontario, Canada. ¹³⁸Hospital Santa Maria, Lisbon, Portugal. ¹³⁹Karadeniz Technical University, Trabzon, Turkey. ¹⁴⁰Istanbul University, Istanbul, Turkey. ¹⁴¹Universidade Federal de Juiz de Fora, Juiz de Fora, Brazil. ¹⁴²Ministry of Public Health, Asunción, Paraguay. ¹⁴³Cardiologia di Mercato S. Severino Hospital, Mercato San Severino, Italy. ¹⁴⁴Karolinska Institutet, Huddinge, Sweden. ¹⁴⁵Centro de Estudios sobre Nutrición Infantil, Buenos Aires, Argentina. ¹⁴⁶University of Porto, Porto, Portugal. ¹⁴⁷Santiago de Compostela University, Santiago de Compostela, Spain. ¹⁴⁸Associazione Calabrese di Epatologia, Reggio Calabria, Italy. ¹⁴⁹Ministry of Health, Ankara, Turkey. ¹⁵⁰Food and Agriculture Organization of the United Nations, Rome, Italy. ¹⁵¹Federal University of Rio Grande, Rio Grande, Brazil. ¹⁵²India Diabetes Research Foundation, Chennai, India. ¹⁵³Duke-NUS Medical School, Singapore, Singapore. ¹⁵⁴National Institute of Medical Statistics, New Delhi, India. ¹⁵⁵Academia Sinica, Taipei, Taiwan. ¹⁵⁶Capital Institute of Pediatrics, Beijing, China. ¹⁵⁷Duke University, Durham, NC, USA. ¹⁵⁸Kailuan General Hospital, Tangshan, China. ¹⁵⁹University of Oxford, Oxford, UK. ¹⁶⁰The Gertner Institute for Epidemiology and Health Policy Research, Ramat Gan, Israel. ¹⁶¹National Centre of Public Health and Analyses, Sofia, Bulgaria. ¹⁶²University of Bern, Lausanne, Switzerland. ¹⁶³Ministry of Health and Welfare, Taipei, Taiwan. ¹⁶⁴Murcia Health Council, Murcia, Spain. ¹⁶⁵Seoul National University College of Medicine, Seoul, South Korea. ¹⁶⁶Korea Centers for Disease Control and Prevention, Cheongju-si, South Korea. ¹⁶⁷Universidade Estadual Paulista, Presidente Prudente, Brazil. ¹⁶⁸Medical University of Silesia, Katowice, Poland. ¹⁶⁹Charles University in Prague, Prague, Czech Republic. ¹⁷⁰Thomayer Hospital, Prague, Czech Republic. ¹⁷¹Primary Health Care, Floriana, Malta. ¹⁷²Carol Davila University of Medicine and Pharmacy, Bucharest, Romania. ¹⁷³Katholieke Universiteit Leuven, Leuven, Belgium. ¹⁷⁴Statistics Canada, Ottawa, Ontario, Canada. ¹⁷⁵Agency for Preventive and Social Medicine, Bregenz, Austria. ¹⁷⁶Universidade Federal de Santa Catarina, Florianópolis, Brazil. ¹⁷⁷University of Southampton, Southampton, UK. ¹⁷⁸Institut Pasteur de Lille, Lille, France. ¹⁷⁹Malawi Epidemiology and Intervention Research Unit, Lilongwe, Malawi. ¹⁸⁰CIBEROBN, Madrid, Spain. ¹⁸¹National Institute of Public Health, Bucharest, Romania. ¹⁸²Eduardo Mondlane University, Maputo, Mozambique. ¹⁸³National Council of Research, Reggio Calabria, Italy. ¹⁸⁴Indian Statistical Institute, Kolkata, India. ¹⁸⁵Tabriz Health Services Management Centre, Tabriz, Iran. ¹⁸⁶Sciensano, Brussels, Belgium. ¹⁸⁷Academic Medical Center of University of Amsterdam, Amsterdam, The Netherlands. ¹⁸⁸National Research Centre for Preventive Medicine, Moscow, Russia. ¹⁸⁹Erasmus Medical Center Rotterdam, Rotterdam, The Netherlands. ¹⁹⁰University of Montreal, Montreal, Québec, Canada. ¹⁹¹Institut de Recherche pour le Développement, Montpellier, France. ¹⁹²French Public Health Agency, St Maurice, France. ¹⁹³Mediterranean Cardiocentro, Naples, Italy. ¹⁹⁴Universidade do Vale do Rio dos Sinos, São Leopoldo, Brazil. ¹⁹⁵National Council of Scientific and Technical Research, Tandil, Argentina. ¹⁹⁶Ministry of Health and Medical Education, Tehran, Iran. ¹⁹⁷National Institute of Nutrition, Hanoi, Vietnam. ¹⁹⁸University of Queensland, Brisbane, Queensland, Australia. ¹⁹⁹Istituto Superiore di Sanità, Rome, Italy. ²⁰⁰Universidad de Cuenca,

Cuenca, Ecuador. ²⁰¹Helmholtz Zentrum München, Munich, Germany. ²⁰²Ministère de la Santé et de la Lutte Contre le Sida, Abidjan, Côte d'Ivoire. ²⁰³The Cardinal Wyszyński Institute of Cardiology, Warsaw, Poland. ²⁰⁴Beijing Center for Disease Prevention and Control, Beijing, China. ²⁰⁵BIDMC, Boston, MA, USA. ²⁰⁶University of Latvia, Riga, Latvia. ²⁰⁷Ministry of Health and Medical Services, Gizo, Solomon Islands. ²⁰⁸University of Benin, Benin City, Nigeria. ²⁰⁹University of Skövde, Skövde, Sweden. ²¹⁰National Institute of Nutrition and Food Technology, Tunis, Tunisia. ²¹¹The University of the West Indies, Kingston, Jamaica. ²¹²University of California Davis, Davis, CA, USA. ²¹³University of Stellenbosch, Cape Town, South Africa. ²¹⁴National Institute for Health and Welfare, Helsinki, Finland. ²¹⁵Queen's University of Belfast, Belfast, UK. ²¹⁶Shiraz University of Medical Sciences, Shiraz, Iran. ²¹⁷Centro de Salud Villanueva Norte, Badajoz, Spain. ²¹⁸Hospital Don Benito-Villanueva de la Serena, Badajoz, Spain. ²¹⁹Ministry of Health, Buenos Aires, Argentina. ²²⁰Council for Agricultural Research and Economics, Rome, Italy. ²²¹Pontificia Universidad Católica de Chile, Santiago, Chile. ²²²Institute of Mother and Child, Warsaw, Poland. ²²³University of Basel, Basel, Switzerland. ²²⁴Swiss TPH, Basel, Switzerland. ²²⁵University of Tartu, Tartu, Estonia. ²²⁶Instituto Nacional de Salud Pública, Cuernavaca, Mexico. ²²⁷Universiti Sains Malaysia, Kelantan, Malaysia. ²²⁸Umeå University, Umeå, Sweden. ²²⁹Georgia College and State University, Milledgeville, GA, USA. ²³⁰Federal University of São Paulo, São Paulo, Brazil. ²³¹Hospital Universitario Son Espases, Palma, Spain. ²³²Hospital de Clínicas de Porto Alegre, Porto Alegre, Brazil. ²³³Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil. ²³⁴Kindai University, Osaka-Sayama, Japan. ²³⁵Kyoto University, Kyoto, Japan. ²³⁶Medical University of Warsaw, Warsaw, Poland. ²³⁷Ministry of Health, Rome, Italy. ²³⁸University of Catania, Catania, Italy. ²³⁹CIBER en Epidemiología y Salud Pública, Alicante, Spain. ²⁴⁰Africa Health Research Institute, Mtubatuba, South Africa. ²⁴¹Geneva University Hospitals, Geneva, Switzerland. ²⁴²CIBER en Epidemiología y Salud Pública, Barcelona, Spain. ²⁴³Australian Bureau of Statistics, Canberra, Australian Capital Territory, Australia. ²⁴⁴Universidade Federal de Minas Gerais, Belo Horizonte, Brazil. ²⁴⁵Wageningen University, Wageningen, The Netherlands. ²⁴⁶University of Insubria, Varese, Italy. ²⁴⁷University of Adelaide, Adelaide, South Australia, Australia. ²⁴⁸Lund University, Lund, Sweden. ²⁴⁹Ministry of Health, Jerusalem, Israel. ²⁵⁰McGill University, Montreal, Québec, Canada. ²⁵¹Universidad Autónoma de Santo Domingo, Santo Domingo, Dominican Republic. ²⁵²Universidad Politécnica de Madrid, Madrid, Spain. ²⁵³The Andes Clinic of Cardio-Metabolic Studies, Merida, Venezuela. ²⁵⁴Instituto Nacional de Higiene, Epidemiología y Microbiología, Havana, Cuba. ²⁵⁵Ministry of Health, Lisbon, Portugal. ²⁵⁶Institute for Clinical and Experimental Medicine, Prague, Czech Republic. ²⁵⁷Children's Memorial Health Institute, Warsaw, Poland. ²⁵⁸Aristotle University of Thessaloniki, Thessaloniki, Greece. ²⁵⁹Jagiellonian University Medical College, Kraków, Poland. ²⁶⁰National Center of Cardiovascular Diseases, Beijing, China. ²⁶¹University of Ferrara, Ferrara, Italy. ²⁶²Icelandic Heart Association, Kopavogur, Iceland. ²⁶³Universidad Icesi, Cali, Colombia. ²⁶⁴State University of Montes Claros, Montes Claros, Brazil. ²⁶⁵King's College London, London, UK. ²⁶⁶International Agency for Research on Cancer, Lyon, France. ²⁶⁷Capital Medical University, Beijing, China. ²⁶⁸Healis-Sekhsaria Institute for Public Health, Navi Mumbai, India. ²⁶⁹Eternal Heart Care Centre and Research Institute, Jaipur, India. ²⁷⁰University of Ibadan, Ibadan, Nigeria. ²⁷¹Institute for Clinical Effectiveness and Health Policy, Buenos Aires, Argentina. ²⁷²Research and Education Institute of Child Health, Nicosia, Cyprus. ²⁷³Danish Cancer Society Research Centre, Copenhagen, Denmark. ²⁷⁴Alexander Technological Educational Institute, Thessaloniki, Greece. ²⁷⁵Kyushu University, Fukuoka, Japan. ²⁷⁶University of Bergen, Bergen, Norway. ²⁷⁷Tulane University, New Orleans, LA, USA. ²⁷⁸Chinese Center for Disease Control and Prevention, Beijing, China. ²⁷⁹University of Pécs, Pécs, Hungary. ²⁸⁰Danish Health Authority, Copenhagen, Denmark. ²⁸¹Joep Lange Institute, Amsterdam, The Netherlands. ²⁸²Oulu University Hospital, Oulu, Finland. ²⁸³Universidad Autónoma de Bucaramanga, Bucaramanga, Colombia. ²⁸⁴ETH Zurich, Zurich, Switzerland. ²⁸⁵Chronic Diseases Research Center, Tehran, Iran. ²⁸⁶University of Hong Kong, Hong Kong, China. ²⁸⁷The Chinese University of Hong Kong, Hong Kong, China. ²⁸⁸University of Western Australia, Perth, Western Australia, Australia. ²⁸⁹Kingston Health Sciences Centre, Kingston, Ontario, Canada. ²⁹⁰Heart Institute, São Paulo, Brazil. ²⁹¹Fundación Oftalmológica de Santander, Santander, Colombia. ²⁹²University Oran 1, Oran, Algeria. ²⁹³Independent Public Health Specialist, Nay Pyi Taw, Myanmar. ²⁹⁴Ministry of Health and Sports, Nay Pyi Taw, Myanmar. ²⁹⁵Peking University, Beijing, China. ²⁹⁶CIBER en Epidemiología y Salud Pública, Murcia, Spain. ²⁹⁷Amsterdam UMC of University of Amsterdam, Amsterdam, The Netherlands. ²⁹⁸Vrije Universiteit Amsterdam, Amsterdam, The Netherlands. ²⁹⁹American University of Beirut, Beirut, Lebanon. ³⁰⁰CIBER en Epidemiología y Salud Pública, San Sebastian, Spain. ³⁰¹Cairo University, Cairo, Egypt. ³⁰²The University of Tokyo, Tokyo, Japan. ³⁰³University Hospital Centre Zagreb, Zagreb, Croatia. ³⁰⁴Niigata University, Niigata, Japan. ³⁰⁵Hadassah University Medical Center, Jerusalem, Israel. ³⁰⁶Norwegian University of Science and Technology, Trondheim, Norway. ³⁰⁷The University of Melbourne, Melbourne, Victoria, Australia. ³⁰⁸Sports University of Tirana, Tirana, Albania. ³⁰⁹University of Zagreb School of Medicine, Zagreb, Croatia. ³¹⁰Heart Foundation, Melbourne, Victoria, Australia. ³¹¹National Health Insurance Service, Wonju, South Korea. ³¹²Guangzhou 12th Hospital, Guangzhou, China. ³¹³Universidad Eugenio María de Hostos, Santo Domingo, Dominican Republic. ³¹⁴Simon Fraser University, Burnaby, British Columbia, Canada. ³¹⁵Ruprecht-Karls-University of Heidelberg, Heidelberg, Germany. ³¹⁶World Health Organization Country Office, Delhi, India. ³¹⁷Institute of Public Health of Serbia, Belgrade, Serbia. ³¹⁸University of Opole, Opole, Poland. ³¹⁹University of Crete, Heraklion, Greece. ³²⁰Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia. ³²¹Hellenic Medical Association for Obesity, Athens, Greece. ³²²Maharajgunj Medical Campus, Kathmandu, Nepal. ³²³Aarhus University, Aarhus, Denmark. ³²⁴Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA. ³²⁵Pennington Biomedical Research Center, Baton Rouge, LA, USA. ³²⁶University of Eastern Finland, Kuopio, Finland. ³²⁷National Institute of Epidemiology, Chennai, India. ³²⁸University of Münster, Münster, Germany. ³²⁹Research Institute for Primordial Prevention of Non-communicable Disease, Isfahan, Iran. ³³⁰Amsterdam Public Health Research Institute, Amsterdam, The Netherlands. ³³¹Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan. ³³²Research Institute of Child Nutrition, Dortmund, Germany. ³³³University of Cambridge, Cambridge, UK. ³³⁴Hypertension Research Center, Isfahan, Iran. ³³⁵Medical University of Innsbruck, Innsbruck, Austria. ³³⁶Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania. ³³⁷National Cancer Center, Goyang-si, South Korea. ³³⁸Statistics Austria, Vienna, Austria. ³³⁹B. P. Koirala Institute of Health Sciences, Dharan, Nepal. ³⁴⁰University of Vienna, Vienna, Austria. ³⁴¹Oulu Deaconess Institute Foundation, Oulu, Finland. ³⁴²Tartu University Clinics, Tartu, Estonia. ³⁴³Kansai Medical University, Osaka-Sayama, Japan. ³⁴⁴National Institute of Pharmacy and Nutrition, Budapest, Hungary. ³⁴⁵Ministry of Health and Quality of Life, Port Louis, Mauritius. ³⁴⁶Polish Academy of Sciences Anthropology Unit, Wrocław, Poland. ³⁴⁷University Hospital Ulm, Ulm, Germany. ³⁴⁸University of Groningen, Groningen, The Netherlands. ³⁴⁹North-West University, Potchefstroom, South Africa. ³⁵⁰National Institute of Public Health, Prague, Czech Republic. ³⁵¹University of Jyväskylä, Jyväskylä, Finland. ³⁵²Institute of Public Health of Montenegro, Podgorica, Montenegro. ³⁵³Amrita Institute of Medical Sciences, Cochin, India. ³⁵⁴Institute of Endocrinology, Prague, Czech Republic. ³⁵⁵All India Institute of Medical Sciences, New Delhi, India. ³⁵⁶African Population and Health Research Center, Nairobi, Kenya. ³⁵⁷Hanoi University of Public Health, Hanoi, Vietnam. ³⁵⁸Higher Institute of Nursing Professions and Technical Health, Rabat, Morocco. ³⁵⁹National Institute of Public Health of Algeria, Algiers, Algeria. ³⁶⁰Croatian National Institute of Public Health, Zagreb, Croatia. ³⁶¹Sahlgrenska Academy, Gothenburg, Sweden. ³⁶²Endocrinology and Metabolism Research Center, Tehran, Iran. ³⁶³University of Public Health, Yangon, Myanmar. ³⁶⁴International Food Policy Research Institute, Dakar, Senegal. ³⁶⁵National University of Singapore, Singapore, Singapore. ³⁶⁶Hong Kong Polytechnic University, Hong Kong, China. ³⁶⁷Tampere University Hospital, Tampere, Finland. ³⁶⁸University of Douala, Douala, Cameroon. ³⁶⁹University of Cape Town, Cape Town, South Africa. ³⁷⁰West Virginia University, Morgantown, WV, USA. ³⁷¹Oswaldo Cruz Foundation Rene Rachou Research Institute, Belo Horizonte, Brazil. ³⁷²National Taiwan University, Taipei, Taiwan. ³⁷³University of Chinese Academy of Sciences, Shanghai, China. ³⁷⁴Uppsala University, Uppsala, Sweden. ³⁷⁵Capital Medical University Beijing An Zhen Hospital, Beijing, China. ³⁷⁶National Institute for Health Development, Tallinn, Estonia. ³⁷⁷Universidad San Martín de Porres, Lima, Peru. ³⁷⁸University Medicine of Greifswald, Greifswald, Germany. ³⁷⁹Consejería de Sanidad Junta de Castilla y León, Valladolid, Spain. ³⁸⁰Norrbottn County Council, Luleå, Sweden. ³⁸¹University of Uppsala, Uppsala, Sweden. ³⁸²Universidade Federal de Ouro Preto, Ouro Preto, Brazil. ³⁸³University of Coimbra, Coimbra, Portugal. ³⁸⁴The Jikei University School of Medicine, Tokyo, Japan. ³⁸⁵National Research Council, Padua, Italy. ³⁸⁶Baker Heart and Diabetes Institute, Melbourne, Victoria, Australia. ³⁸⁷Agricultural University of Athens, Athens, Greece. ³⁸⁸Hospital Israelita Albert Einstein, São Paulo, Brazil. ³⁸⁹Institute of Internal and Preventive Medicine, Novosibirsk, Russia. ³⁹⁰Harokopio University, Athens, Greece. ³⁹¹University of Otago, Dunedin, New Zealand. ³⁹²University of Padua, Padua, Italy. ³⁹³Technological Educational Institute of Crete, Heraklion, Greece. ³⁹⁴Loughborough University, Loughborough, UK. ³⁹⁵Ministry of Health, Nicosia, Cyprus. ³⁹⁶Lausanne University Hospital, Lausanne, Switzerland. ³⁹⁷CIBERCV, Barcelona, Spain. ³⁹⁸Mary Immaculate College, Limerick, Ireland. ³⁹⁹Emory University, Atlanta, GA, USA. ⁴⁰⁰Hungarian Society of Sports Medicine, Budapest, Hungary. ⁴⁰¹Sher-i-Kashmir Institute of Medical Sciences, Srinagar, India. ⁴⁰²UiT The Arctic University of Norway, Tromsø, Norway. ⁴⁰³National Centre for Disease Informatics and Research, New Delhi, India. ⁴⁰⁴Cape Peninsula University of Technology, Cape Town, South Africa. ⁴⁰⁵University of Rzeszów, Rzeszów, Poland. ⁴⁰⁶University of Edinburgh, Edinburgh, UK. ⁴⁰⁷International Medical University, Shah Alam, Malaysia. ⁴⁰⁸Institut National de la Santé et de la Recherche Médicale, Lille, France. ⁴⁰⁹Robert Koch Institute, Berlin, Germany. ⁴¹⁰Lusófona University, Lisbon, Portugal. ⁴¹¹Democritus University, Alexandroupolis, Greece. ⁴¹²Grigore T. Popa University of Medicine and Pharmacy, Iasi, Romania. ⁴¹³Università degli Studi di Firenze, Florence, Italy. ⁴¹⁴Isfahan Cardiovascular Research Center, Isfahan, Iran. ⁴¹⁵Strasbourg University Hospital, Strasbourg, France. ⁴¹⁶University of Strasbourg, Strasbourg, France. ⁴¹⁷Mulago Hospital, Kampala, Uganda. ⁴¹⁸Instituto Nacional de Salud Pública, Mexico City, Mexico. ⁴¹⁹University of Limpopo, Sovenga, South Africa. ⁴²⁰Seoul National University Children's Hospital, Seoul, South Korea. ⁴²¹University Medical Science, Havana, Cuba. ⁴²²Universidad de Zaragoza, Zaragoza, Spain. ⁴²³RCSI, Dublin, Ireland. ⁴²⁴La Trobe University, Melbourne, Victoria, Australia. ⁴²⁵International Institute of Molecular and Cell Biology, Warsaw, Poland. ⁴²⁶Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. ⁴²⁷Gorgas Memorial Institute of Public Health, Panama City, Panama. ⁴²⁸World Health Organization Country Office, Lilongwe, Malawi. ⁴²⁹Department of Public Health, Nay Pyi Taw, Myanmar. ⁴³⁰University of Brescia, Brescia, Italy. ⁴³¹Ministry of Health and Social Protection, Dushanbe, Tajikistan. ⁴³²Bushehr University of Medical Sciences, Bushehr, Iran. ⁴³³Ulm University, Ulm, Germany. ⁴³⁴Kobe University, Kobe, Japan. ⁴³⁵Suraj Eye Institute, Nagpur, India. ⁴³⁶UNICEF, Yaoundé, Cameroon. ⁴³⁷National Institute of Hygiene and Epidemiology, Hanoi, Vietnam. ⁴³⁸University of Pharmacy and Medicine, Ho Chi Minh City, Vietnam. ⁴³⁹Hanoi Medical University, Hanoi, Vietnam. ⁴⁴⁰Miami Veterans Affairs Healthcare System, Miami, FL, USA. ⁴⁴¹Heartfile, Islamabad, Pakistan. ⁴⁴²National Cancer Center, Tokyo, Japan. ⁴⁴³Eastern Mediterranean Public Health Network, Amman, Jordan. ⁴⁴⁴State University of Medicine and Pharmacy, Chisinau, Moldova. ⁴⁴⁵Tachikawa General Hospital, Nagaoka, Japan. ⁴⁴⁶Japan Wildlife Research Center, Tokyo, Japan. ⁴⁴⁷University of Vale do Rio dos Sinos, São Leopoldo, Brazil. ⁴⁴⁸National Food and Nutrition Institute, Warsaw, Poland. ⁴⁴⁹University of Manchester, Manchester, UK. ⁴⁵⁰Ministry of Health, Bandar Seri Begawan, Brunei Darussalam. ⁴⁵¹University of Madeira, Funchal, Portugal. ⁴⁵²University of Puerto Rico, San Juan, Puerto Rico. ⁴⁵³Research Center for Prevention and Health, Glostrup, Denmark. ⁴⁵⁴MRC Lifecourse Epidemiology Unit, Southampton, UK. ⁴⁵⁵University of Novi Sad, Novi Sad, Serbia. ⁴⁵⁶Kwame Nkrumah University of Science and Technology, Kumasi, Ghana. ⁴⁵⁷Institute for Social and Preventive Medicine, Lausanne, Switzerland. ⁴⁵⁸Cancer Prevention and Research Institute, Florence, Italy. ⁴⁵⁹University of Wisconsin-Madison, Madison, WI, USA. ⁴⁶⁰IRCCS Casa Sollievo della Sofferenza, Bari, Italy. ⁴⁶¹Zayed University, Abu Dhabi, United Arab Emirates. ⁴⁶²Catholic University of Daegu, Daegu, South Korea. ⁴⁶³University of Medicine, Pharmacy, Science and Technology of Târgu Mureș, Târgu Mureș, Romania. ⁴⁶⁴Jivandeep Hospital, Anand, India. ⁴⁶⁵South African Medical Research Council, Durban, South Africa. ⁴⁶⁶Spanish Food Safety and Nutrition Agency, Madrid, Spain. ⁴⁶⁷Vietnam National Heart Institute, Hanoi, Vietnam. ⁴⁶⁸Leibniz Institute for Prevention Research and Epidemiology - BIPS, Bremen, Germany. ⁴⁶⁹University of Sarajevo, Sarajevo, Bosnia and Herzegovina. ⁴⁷⁰Cardiovascular Prevention Centre Udine, Udine, Italy. ⁴⁷¹University Hospital of Pisa, Pisa, Italy. ⁴⁷²Ministry of Health and Medical Services, Honiara, Solomon Islands. ⁴⁷³Public Health Agency of Catalonia, Barcelona, Spain. ⁴⁷⁴Institut

Hospital del Mar d'Investigacions Mèdiques, Barcelona, Spain. ⁴⁷⁵Digestive Oncology Research Center, Tehran, Iran. ⁴⁷⁶Digestive Disease Research Institute, Tehran, Iran. ⁴⁷⁷Centre for Disease Prevention and Control, Riga, Latvia. ⁴⁷⁸Alborz University of Medical Sciences, Karaj, Iran. ⁴⁷⁹Ministry of Health, Hanoi, Vietnam. ⁴⁸⁰BRAC, Dhaka, Bangladesh. ⁴⁸¹Institute of Epidemiology Disease Control and Research, Dhaka, Bangladesh. ⁴⁸²University of Turku, Turku, Finland. ⁴⁸³Institut Universitari d'Investigació en Atenció Primària Jordi Gol, Girona, Spain. ⁴⁸⁴Universiti Putra Malaysia, Serdang, Malaysia. ⁴⁸⁵University of Malaya, Kuala Lumpur, Malaysia. ⁴⁸⁶National Institute of Public Health, Copenhagen, Denmark. ⁴⁸⁷University of Valencia, Valencia, Spain. ⁴⁸⁸University of the Philippines, Manila, The Philippines. ⁴⁸⁹Slovak Academy of Sciences, Bratislava, Slovakia. ⁴⁹⁰Nutrition Research Foundation, Barcelona, Spain. ⁴⁹¹Minas Gerais State Secretariat for Health, Belo Horizonte, Brazil. ⁴⁹²Health Center San Agustín, Palma, Spain. ⁴⁹³PharmAccess Foundation, Amsterdam, The Netherlands. ⁴⁹⁴National Institute of Health Doutor Ricardo Jorge, Lisbon, Portugal. ⁴⁹⁵Universidade Nove de Julho, São Paulo, Brazil. ⁴⁹⁶Public Health Agency of Canada, Ottawa, Ontario, Canada. ⁴⁹⁷Canarian Health Service, Tenerife, Spain. ⁴⁹⁸Universidad Industrial de Santander, Santander, Colombia. ⁴⁹⁹Sahlgrenska University Hospital, Gothenburg, Sweden. ⁵⁰⁰Fiji National University, Suva, Fiji. ⁵⁰¹Spanish Nutrition Foundation, Madrid, Spain. ⁵⁰²Institute of Food Sciences of the National Research Council, Avellino, Italy. ⁵⁰³Singapore Eye Research Institute, Singapore, Singapore. ⁵⁰⁴Sitaram Bhartia Institute of Science and Research, New Delhi, India. ⁵⁰⁵Maragheh University of Medical Sciences, Maragheh, Iran. ⁵⁰⁶University of Helsinki, Helsinki, Finland. ⁵⁰⁷National Institute of Health, Lima, Peru. ⁵⁰⁸Ministry of Health, Jakarta, Indonesia. ⁵⁰⁹Catalan Department of Health, Barcelona, Spain. ⁵¹⁰Biodonostia Health Research Institute, San Sebastian, Spain. ⁵¹¹Universidade de Lisboa, Lisbon, Portugal. ⁵¹²South Karelia Social and Health Care District, Lappeenranta, Finland. ⁵¹³Cardiovascular Research Institute, Isfahan, Iran. ⁵¹⁴University of São Paulo Clinics Hospital, São Paulo, Brazil. ⁵¹⁵Hospital Italiano de Buenos Aires, Buenos Aires, Argentina. ⁵¹⁶Medical University of Vienna, Vienna, Austria. ⁵¹⁷Rigshospitalet, Copenhagen, Denmark. ⁵¹⁸Lagos State University College of Medicine, Lagos, Nigeria. ⁵¹⁹University of Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain. ⁵²⁰National Center for Disease Control and Public Health, Tbilisi, Georgia. ⁵²¹National Center for Global Health and Medicine, Tokyo, Japan. ⁵²²Samsung Medical Center, Seoul, South Korea. ⁵²³St Vincent's Hospital, Sydney, New South Wales, Australia. ⁵²⁴University of New South Wales, Sydney, New South Wales, Australia. ⁵²⁵Health Polytechnic Jakarta II Institute, Jakarta, Indonesia. ⁵²⁶Diponegoro University, Semarang, Indonesia. ⁵²⁷University of Bari, Bari, Italy. ⁵²⁸Institut Régional de Santé Publique, Ouidah, Benin. ⁵²⁹University of Bordeaux, Bordeaux, France. ⁵³⁰Institute of Public Health, Skopje, Macedonia. ⁵³¹University of Leuven, Leuven, Belgium. ⁵³²Lamprecht und Stamm Sozialforschung und Beratung AG, Zurich, Switzerland. ⁵³³INSERM, Nancy, France. ⁵³⁴Bonn University, Bonn, Germany. ⁵³⁵Sotiria Hospital, Sotiria, Greece. ⁵³⁶National Institute of Public Health-National Institute of Hygiene, Warsaw, Poland. ⁵³⁷Swansea University, Swansea, UK. ⁵³⁸Fu Jen Catholic University, Taipei, Taiwan. ⁵³⁹National Statistic Office of Cabo Verde, Praia, Cabo Verde. ⁵⁴⁰University of KwaZulu-Natal, Mtubatuba, South Africa. ⁵⁴¹Ministry of Health, Amman, Jordan. ⁵⁴²Comenius University, Bratislava, Slovakia. ⁵⁴³Health Service of Murcia, Murcia, Spain. ⁵⁴⁴IB-SALUT Area de Salut de Menorca, Maó, Spain. ⁵⁴⁵University of Bologna, Bologna, Italy. ⁵⁴⁶Hellenic Health Foundation, Athens, Greece. ⁵⁴⁷Government Medical College, Bhavnagar, India. ⁵⁴⁸Sefako Makgatho Health Science University, Ga-Rankuwa, South Africa. ⁵⁴⁹Addis Ababa University, Addis Ababa, Ethiopia. ⁵⁵⁰Dasman Diabetes Institute, Kuwait City, Kuwait. ⁵⁵¹Ministry of Health, Wellington, New Zealand. ⁵⁵²Universidad Centro-Occidental Lisandro Alvarado, Barquisimeto, Venezuela. ⁵⁵³Meharry Medical College, Nashville, TN, USA. ⁵⁵⁴Dokuz Eylul University, Izmir, Turkey. ⁵⁵⁵University of Tampere Tays Eye Center, Tampere, Finland. ⁵⁵⁶Polytechnic Institute of Porto, Porto, Portugal. ⁵⁵⁷Utrecht University, Utrecht, The Netherlands. ⁵⁵⁸University Medical Center Utrecht, Utrecht, The Netherlands. ⁵⁵⁹National Research Council, Pisa, Italy. ⁵⁶⁰Universidad Miguel Hernandez, Alicante, Spain. ⁵⁶¹Ministry of Health, Mont Fleuri, Seychelles. ⁵⁶²North Karelian Center for Public Health, Joensuu, Finland. ⁵⁶³University of the Witwatersrand, Johannesburg, South Africa. ⁵⁶⁴Institute for Medical Research, Kuala Lumpur, Malaysia. ⁵⁶⁵Xinjiang Medical University, Urumqi, China. ⁵⁶⁶Shanghai Educational Development Co. Ltd, Shanghai, China. ⁵⁶⁷Paracelsus Medical University, Salzburg, Austria. ⁵⁶⁸St George's, University of London, London, UK. ⁵⁶⁹Universitas Indonesia, Jakarta, Indonesia. ⁵⁷⁰Institute of Food and Nutrition Development of Ministry of Agriculture, Beijing, China. ⁵⁷¹Children's Hospital of Fudan University, Shanghai, China. ⁵⁷²University of Cyprus, Nicosia, Cyprus. ⁵⁷³Iran University of Medical Sciences, Tehran, Iran. ⁵⁷⁴West Kazakhstan State Medical University, Aktobe, Kazakhstan. ⁵⁷⁵Inner Mongolia Medical University, Hohhot, China. ⁵⁷⁶Deceased: Deepak N. Amarapurkar, Konrad Jamrozik, Dimitrios Trichopoulos. *e-mail: majid.ezzati@imperial.ac.uk

METHODS

Our aim was to estimate trends in mean BMI from 1985 to 2017 by rural and urban place of residence for 200 countries and territories (Supplementary Table 2). To achieve this aim, we pooled cross-sectional population-based data on height and weight in adults aged 18 years and older. Therefore, by design, our results measure total change in BMI in each country's rural and urban populations, which consists of (1) change in the BMI of individuals due to change in their economic status and environment, and (2) change in the composition of individuals that make up the population (and their economic status and environment). Change in population composition occurs naturally owing to fertility and mortality, as well as owing to migration. Therefore, our results should not be interpreted as solely a change in the BMI of individuals. Both components of change are relevant for policy formulation because policies should address the environment and nutrition of the contemporary population.

We used mean BMI as the primary outcome, rather than prevalence of overweight or obesity, because the relationship between BMI and disease risk is continuous, with each unit lower BMI being associated with a constant proportional reduction in disease risk until a BMI of around 21–23 kg m⁻², which is below the cut-offs used to define overweight and obesity^{37–39}. Therefore, the largest health benefits of weight management are achieved by lowering the population distribution of BMI. Mean BMI is the simplest summary statistic of the population distribution. Nonetheless, mean BMI and prevalence of overweight and obesity are closely associated (Extended Data Fig. 5).

Data sources. We used a database on cardiometabolic risk factors collated by the Non-Communicable Disease Risk Factor Collaboration (NCD-RisC). NCD-RisC is a worldwide network of health researchers and practitioners, that systematically documents the worldwide trends and variations in risk factors for non-communicable diseases. The database was collated through multiple routes for identifying and accessing data. We accessed publicly available population-based measurement surveys—for example, Demographic and Health Surveys, Global School-based Student Health Surveys, the European Health Interview and Health Examination Surveys and those available via the Inter-University Consortium for Political and Social Research. We requested, through the World Health Organization (WHO) and its regional and country offices, help with identification and access to population-based surveys from ministries of health and other national health and statistical agencies. Requests were also sent by the World Heart Federation to its national partners. We made similar requests to the co-authors of an earlier pooled analysis of cardiometabolic risk factors^{40–43} and invited them to reanalyse data from their studies and join NCD-RisC. Finally, to identify major sources not accessed through the above routes, we searched and reviewed published studies as described previously⁴⁴ and invited all eligible studies to join NCD-RisC.

Anonymized individual record data from sources included in NCD-RisC were reanalysed according to a common protocol. Within each survey, we included participants aged 18 years and older who were not pregnant. We dropped participants with implausible BMI levels (defined as BMI < 10 kg m⁻² or BMI > 80 kg m⁻²) or with implausible height or weight values (defined as height < 100 cm, height > 250 cm, weight < 12 kg or weight > 300 kg; <0.2% of all subjects). We also dropped participants whose urban and rural status was unknown in surveys that had recorded place of residence (0.05% of all participants). We calculated mean BMI and its standard error by sex, age group (18 years, 19 years, 10-year age groups from 20–29 years to 70–79 years and 80+ years) and rural or urban place of residence. All analyses incorporated appropriate sample weights and complex survey design, when applicable, in calculating summary statistics. Countries typically use the rural and urban classification of communities designated by their statistical offices at any given time both for survey design and for reporting of population to the United Nations Population Division. The classification can change, for example as previously rural areas grow and industrialize and hence become, and are (re)designated as, *de novo* cities. To the extent that the reclassifications keep up with changes in the real status of each community, survey and population data reflect the status of each community at the time of measurement. For surveys without information on place of residence, we calculated age- and sex-stratified summary statistics for the entire sample, which represented the population-weighted sum of rural and urban means.

To ensure summaries were prepared according to the study protocol, computer code was provided to NCD-RisC members who requested assistance. All submitted data were checked by at least two independent reviewers. Questions and clarifications were discussed with NCD-RisC members and resolved before data were incorporated into the database.

Finally, we incorporated all nationally representative data from sources that were identified but not accessed through the above routes, by extracting summary statistics from published reports. Data were also extracted for nine WHO STEPwise approach to Surveillance (STEPS) surveys, one Countrywide Integrated Non-communicable Diseases Intervention (CINDI) survey, and five sites of the

WHO Multinational MONItoring of trends and determinants in Cardiovascular disease (MONICA) project that were not deposited in the MONICA Data Centre. Data were extracted from published reports only when reported by sex and in age groups no wider than 20 years. We also used data from a previous global data pooling study⁴⁵ when such data had not been accessed through the routes described.

All NCD-RisC members are asked periodically to review the list of sources from their country, to suggest additional sources not in the database, and to verify that the included data meet the inclusion criteria listed below and are not duplicates. The NCD-RisC database is continuously updated through this contact with NCD-RisC members. For this paper, we used data from the NCD-RisC database for years 1985 to 2017 and ages 18 years and older. A list of the data sources that we used in this analysis and their characteristics is provided in Supplementary Table 1.

Data inclusion and exclusion. Data sources were included in the NCD-RisC database if: (1) measured data on height, weight, waist circumference or hip circumference were available; (2) study participants were 5 years of age and older; (3) data were collected using a probabilistic sampling method with a defined sampling frame; (4) data were from population samples at the national, sub-national (that is, covering one or more sub-national regions, more than three urban communities or more than five rural communities) or community level; and (5) data were from the countries and territories listed in Supplementary Table 2.

We excluded all data sources that were based solely on self-reported weight and height without a measurement component, because these data are subject to biases that vary by geography, time, age, sex and socioeconomic characteristics^{45–47}. Owing to these variations, approaches to correcting self-reported data leave residual bias. We also excluded data sources on population subgroups whose anthropometric status may differ systematically from the general population, including: (1) studies that included or excluded people based on their health status or cardiovascular risk; (2) studies whose participants were only ethnic minorities; (3) specific educational, occupational, or socioeconomic subgroups, with the exception noted below; (4) those recruited through health facilities, with the exception noted below; and (5) women aged 15–19 years in surveys which sampled only ever-married women or measured height and weight only among mothers.

We used school-based data in countries, and in age–sex groups, with school enrolment of 70% or higher. We used data for which the sampling frame was health insurance schemes in countries in which at least 80% of the population were insured. Finally, we used data collected through general practice and primary care systems in high-income and central European countries with universal insurance, because contact with the primary care systems tends to be as good as or better than response rates for population-based surveys.

Conversion of BMI prevalence metrics to mean BMI. In 2% of our data points—mostly extracted from published reports or from a previous pooling analysis⁴³—mean BMI was not reported, but data were available for the prevalence of one or more BMI categories, for example, BMI ≥ 30 kg m⁻². In order to use these data, we used previously validated conversion regressions² to estimate the missing primary outcome from the available BMI prevalence metric(s). All sources of uncertainty in the conversion—including the sampling uncertainty of the original data, the uncertainty of the regression coefficients and random effects, and the regression residuals—were carried forward by using repeated draws from their joint posterior distribution, accounting for the correlations among the uncertainties of regression coefficients and random effects.

Statistical analysis of BMI trends by rural and urban place of residence. We used a Bayesian hierarchical model to estimate mean BMI by country, year, sex, age and place of residence. The statistical model is described in detail in a statistical paper and related substantive papers^{2,35,40–44,48–51}, and in the Supplementary Information. In summary, we organized countries into 21 regions (Supplementary Table 2), mostly based on geography and national income. The exception was high-income English-speaking countries (Australia, Canada, Ireland, New Zealand, the United Kingdom and the United States), grouped together in one region because BMI and other cardiometabolic risk factors have similar trends in these countries, which can be distinct from other countries in their geographical regions^{2,49,50,52}. Regions were in turn organized into nine super-regions.

The model had a hierarchical structure in which estimates for each country and year were informed by their own data, if available, and by data from other years in the same country and from other countries, especially those in the same region with data for similar time periods. The extent to which estimates for each country-year were influenced by data from other years and other countries depended on whether the country had data, the sample size of the data, whether they were national, and the within-country and within-region variability of the available data. The model incorporated nonlinear time trends comprising linear terms and a second-order random walk, all modelled hierarchically. The age association of BMI was modelled using a cubic spline to allow nonlinear age patterns, which could vary across countries. The model accounted for the possibility that BMI in sub-national and community samples might differ systematically from nationally

representative ones and have larger variation than in national studies. These features were implemented by including data-driven fixed-effect and random-effect terms for sub-national and community data. The fixed effects adjusted for systematic differences between sub-national or community studies and national studies. The random effects allowed national data to have larger influence on the estimates than sub-national or community data with similar sample sizes.

Here, we extended the model to make estimates for rural and urban populations following a previously published approach^{35,51}. This model includes a parameter representing the urban–rural BMI difference, which is estimated empirically and allowed to vary by country and year. The model uses all of the data—those stratified by rural and urban place of residence as well as those reported for the entire population. If data for a country–year were not stratified by place of residence, the estimated urban–rural BMI difference was informed by stratified data from other years and countries, especially those in the same region with data from similar time periods.

We fitted the statistical model with the Markov chain Monte Carlo (MCMC) algorithm and following burn-in obtained 5,000 samples (or draws) from the posterior distribution of model parameters, which were in turn used to obtain the posterior distributions of our primary outcomes—mean urban BMI, mean rural BMI and mean urban–rural BMI difference. Posterior estimates were made in 1-year age groups for ages 18 and 19 and 5-year age groups for those aged 20 years and older. We generated age-standardized estimates by taking weighted means of age-specific estimates, using age weights from the WHO standard population. Regional and global rural and urban mean BMI estimates were calculated as population-weighted averages of rural and urban mean for the constituent country estimates by age group and sex. National mean BMI was calculated as population-weighted averages of the rural and urban means. All analyses were done separately by sex because geographical and temporal patterns of BMI differ between men and women².

The reported credible intervals represent the 2.5th and the 97.5th percentiles of the posterior distributions. We report the posterior probability that the estimated urban–rural BMI difference is a true difference in the same direction as the posterior mean estimate. We also report the posterior probability that the estimated change in the rural–urban BMI difference over time represents a true increase or decrease.

Validation of statistical model. We calculated the difference between the posterior estimates from the model and data from national studies. Median errors were very close to zero (0.03 kg m⁻² for women and -0.02 kg m⁻² for men) and median absolute errors were 0.32 kg m⁻² for women and 0.26 kg m⁻² for men, indicating that the estimates were unbiased and had small deviations relative to national studies. The differences were indistinguishable from zero at the 5% level of statistical significance.

We also tested how well our statistical model predicts missing data, known as external predictive validity or cross-validation, in two different tests. In the first test, we held out all data from 10% of countries with data (that is, created the appearance of countries with no data for which we actually had data). The countries for which the data were withheld were selected randomly from the following three groups: data rich (8 or more data sources for women and 7 or more data sources for men), data poor (1–3 data sources for women and 1–2 for men) and average data availability (4–7 data sources for women and 3–6 for men). All data-rich countries had at least one data source after 2000 and at least one source with data stratified on rural and urban place of residence. We fitted the model to the data from the remaining 90% of countries and made estimates of the held-out observations. In the second test, we assessed other patterns of missing data by holding out 10% of our data sources, again from a mix of data-rich, data-poor and average-data countries, as defined above. For a given country, we either held out a random one third of the country's data or all of the country's 2000–2017 data to determine, respectively, how well we filled in the gaps for countries with intermittent data and how well we estimated in countries without recent data. We fitted the model to the remaining 90% of the dataset and made estimates of the held-out observations. We repeated each test five times, holding out a different subset of data in each repetition. In both tests, we calculated the differences between the held-out data and the estimates. We also calculated the 95% credible intervals of the estimates; in a model with good external predictive validity, 95% of held-out values would be included in the 95% credible intervals.

Our statistical model performed very well in the external validation tests, that is, in estimating mean BMI when data were missing. The estimates of mean BMI were unbiased, as evidenced with median errors that were zero or close to zero globally (0.03 and -0.03 kg m⁻² for women and -0.15 and 0.00 kg m⁻² for men in the first and second tests, respectively), and less than ± 0.20 kg m⁻² in every subset of withheld data except 1985–1999 data in the first test for men, for which the median error was -0.24 kg m⁻² (Extended Data Table 2). Most of the median errors were indistinguishable from zero at the 5% level of statistical significance.

The 95% credible intervals of estimated mean BMI covered 94–98% of true data globally; coverage was >93% in all but one subset of withheld data. Median absolute errors ranged from 0.52 to 1.09 kg m⁻² globally and were at most 1.29 kg m⁻² in all subsets of withheld data. Median absolute errors were smaller in the second test, in which subsets of data sources from some countries were withheld, than in the first test, in which all data from some countries were withheld. Given that we had data for 190 out of 200 countries for women and 183 out of 200 countries for men, the second test is a better reflection of data availability in our analysis. For comparison, median absolute differences for mean BMI between pairs of nationally representative surveys done in the same country and in the same year was 0.46 kg m⁻², indicating that our estimates perform almost as well as running two parallel surveys in the same country and year.

Contributions of urbanization and rural and urban BMI change to changes in population mean BMI. We calculated the contributions of the following components to change in population mean BMI from 1985 to 2017: the contribution of change in BMI in rural areas, the contribution of change in BMI in urban areas, and the contribution of urbanization (that is, increase in the proportion of people living in urban areas). The first two parts were calculated by fixing the proportion of people living in rural and urban areas to 1985 levels and allowing BMI to change as it did in the respective population. The contribution of urbanization was calculated by fixing BMI in rural and urban areas to 2017 levels and allowing the proportion of people living in cities to change as it did. Percentage contributions were calculated using posterior draws, with reported credible intervals representing the 2.5th and the 97.5th percentiles of their posterior distributions. The change in mean BMI from 1985 to 2017 was then calculated as (contribution of change in rural BMI + contribution of change in urban BMI + contribution of change in the proportion of the population living in urban areas) = ((change in BMI_{rural1985–2017}) (percentage living in rural areas₁₉₈₅) + (change in BMI_{urban1985–2017}) (percentage living in urban areas₁₉₈₅) + (change in percentage living in urban areas_{1985–2017}) (BMI_{urban2017} - BMI_{rural2017})).

Strengths and limitations. Urbanization is regarded as one of the most important contributors to the global obesity epidemic, but this perspective is based on limited data. We present the first comparable estimates of mean BMI for rural and urban populations worldwide over three decades using, to our knowledge, the largest and most comprehensive global database of human anthropometry with information on urban or rural place of residence. We used population-based measurement data from almost all countries, with information on participants' urban or rural place of residence for the majority of data sources. We maintained a high level of data quality through repeated checks of study characteristics against our inclusion and exclusion criteria, which were verified by NCD-RisC members, and did not use any self-reported data to avoid bias in height and weight. Data were analysed according to a common protocol to obtain mean BMI by age, sex and place of residence. We used a statistical model that used all available data, while giving more weight to national data than sub-national and community studies and took into account the epidemiological features of BMI by using nonlinear time trends and age associations. The model used information on the urban–rural difference in BMI where available and estimated this difference hierarchically and temporally in the absence of stratified data.

Despite our large-scale data collation effort, some countries and regions had fewer data sources, particularly the Caribbean, and Polynesia and Micronesia. There were also fewer data sources before 2000. This temporal and geographical sparsity of data led to wider uncertainty intervals for these countries, regions and years. Although health surveys commonly use the rural and urban classification of national statistical offices, cities and rural areas in different countries vary in their demographic characteristics (for example, population size or density), economic activity, administrative structures, infrastructure and environment. These differences appropriately exist because countries themselves differ in terms of their demography, geography and economy. For example, a country with a smaller population may use a lower threshold for urban designation than one with a larger population, because its cities are naturally smaller even if they serve the same functions. Official rural and urban classifications are used for resource allocation and planning for nutrition and health^{53–58}, which makes them the appropriate unit for tracking outcomes. Nonetheless, understanding the causes of change in rural and urban areas can be enriched with use of more complex and multi-dimensional measures of urbanicity involving size, density, economic and commercial activities and infrastructures^{59,60}. Finally, urbanization could arise from a variety of mechanisms: (1) natural increase due to excess births over deaths in cities compared to rural areas, (2) rural to urban migration (often related to opportunities for work and education) and (3) reclassification of previously rural areas as they grow and industrialize and hence become, and are (re)designated as, *de novo* cities. The contributions of these mechanisms to urbanization vary across countries. The use of time-varying rural versus urban classification of communities ensures that in any year, the rural and urban strata represent the actual status of each community.

However, each of these mechanisms may have different implications for changes in nutrition and physical activity and, therefore, BMI.


Data availability

Estimates of mean BMI by country, year, sex and urban and rural place of residence are available from <http://www.ncdrisc.org/>. Input data from publicly available sources can also be downloaded from <http://www.ncdrisc.org/>. For other data sources, contact information for data providers can be obtained from <http://www.ncdrisc.org/>.


Code availability

The computer code for the Bayesian hierarchical model used in this work is available at <http://www.ncdrisc.org/>.


37. Prospective Studies Collaboration. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* **373**, 1083–1096 (2009).
38. The Global BMI Mortality Collaboration. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet* **388**, 776–786 (2016).
39. Singh, G. M. et al. The age-specific quantitative effects of metabolic risk factors on cardiovascular diseases and diabetes: a pooled analysis. *PLoS ONE* **8**, e65174 (2013).
40. Danaei, G. et al. National, regional, and global trends in systolic blood pressure since 1980: systematic analysis of health examination surveys and epidemiological studies with 786 country-years and 5.4 million participants. *Lancet* **377**, 568–577 (2011).
41. Danaei, G. et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet* **378**, 31–40 (2011).
42. Farzadfar, F. et al. National, regional, and global trends in serum total cholesterol since 1980: systematic analysis of health examination surveys and epidemiological studies with 321 country-years and 3.0 million participants. *Lancet* **377**, 578–586 (2011).
43. Finucane, M. M. et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* **377**, 557–567 (2011).
44. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* **387**, 1377–1396 (2016).
45. Hayes, A. J., Clarke, P. M. & Lung, T. W. C. Change in bias in self-reported body mass index in Australia between 1995 and 2008 and the evaluation of correction equations. *Popul. Health Metr.* **9**, 53 (2011).
46. Gorber, S. C., Tremblay, M., Moher, D. & Gorber, B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes. Rev.* **8**, 307–326 (2007).
47. Ezzati, M., Martin, H., Skjold, S., Vander Hoorn, S. & Murray, C. J. L. Trends in national and state-level obesity in the USA after correction for self-report bias: analysis of health surveys. *J. R. Soc. Med.* **99**, 250–257 (2006).
48. Finucane, M. M., Paciorek, C. J., Danaei, G. & Ezzati, M. Bayesian estimation of population-level trends in measures of health status. *Stat. Sci.* **29**, 18–25 (2014).
49. NCD Risk Factor Collaboration (NCD-RisC). A century of trends in adult human height. *eLife* **5**, e13410 (2016).
50. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet* **389**, 37–55 (2017).
51. Finucane, M. M., Paciorek, C. J., Stevens, G. A. & Ezzati, M. Semiparametric Bayesian density estimation with disparate data sources: a meta-analysis of global childhood undernutrition. *J. Am. Stat. Assoc.* **110**, 889–901 (2015).
52. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* **387**, 1513–1530 (2016).
53. The World Bank. *An Overview of Links Between Obesity and Food Systems: Implications for the Food and Agriculture Global Practice Agenda*. <http://documents.worldbank.org/curated/en/222101499437276873/pdf/117200REVISED-WP-Obesity-Overview-Web-PUBLIC-002.pdf> (World Bank, 2017).
54. FAO. *Ending Poverty and Hunger by Investing in Agriculture and Rural Areas*. <http://www.fao.org/3/a-i7556e.pdf> (Food and Agriculture Organization of the United Nations, 2017).
55. WHO & Global Health Workforce Alliance. *Global Experience of Community Health Workers for Delivery of Health Related Millenium Development Goals: a Systematic Review, Country Case Studies, and Recommendations for Integration into National Health Systems*. https://www.who.int/workforcealliance/knowledge/publications/CHW_FullReport_2010.pdf?ua=1 (World Health Organization, 2010).
56. Bhutta, Z. A. et al. Rethinking community based strategies to tackle health inequities in South Asia. *Br. Med. J.* **363**, k4884 (2018).
57. IFAD. *Rural Development Report 2016: Fostering Inclusive Rural Transformation*. <https://www.ifad.org/documents/38714170/39155702/Rural+development+report+2016.pdf/347402dd-a37f-41b7-9990-aa745dc113b9> (International Fund for Agricultural Development, 2016).
58. FAO, IFAD & WFP. *The State of Food Insecurity in the World 2015. Meeting the 2015 International Hunger Targets: Taking Stock of Uneven Progress*. <http://www.fao.org/3/a-i4646e.pdf> (Food and Agriculture Organization of the United Nations, 2015).
59. Dahly, D. L. & Adair, L. S. Quantifying the urban environment: a scale measure of urbanicity outperforms the urban–rural dichotomy. *Soc. Sci. Med.* **64**, 1407–1419 (2007).
60. Jones-Smith, J. C. & Popkin, B. M. Understanding community context and adult health changes in China: development of an urbanicity scale. *Soc. Sci. Med.* **71**, 1436–1446 (2010).




Extended Data Fig. 1 | Number of data sources by country. The colour indicates the number of population-based data sources used in the analysis for each country. Countries and territories not included in the analysis are coloured in grey.



Extended Data Fig. 2 | Age-standardized national, rural and urban mean BMI in women aged 18 years and older in 1985 and 2017 by country. The numerical values are provided in Supplementary Table 3 and can be downloaded from <http://www.ncdrisc.org>.




Extended Data Fig. 3 | Age-standardized national, rural and urban mean BMI in men aged 18 years and older in 1985 and 2017 by country. The numerical values are provided in Supplementary Table 3 and can be downloaded from <http://www.ncdrisc.org>.




Extended Data Fig. 4 | The difference between rural and urban age-standardized mean BMI in 1985 compared to 2017. Each point shows one country and colours indicate region. A positive number indicates a higher urban mean BMI and a negative number indicates a higher rural mean BMI. Different sections labelled A–F indicate the following categories of countries. A, countries with an urban excess BMI that

increased from 1985 to 2017. B, countries with an urban excess BMI that decreased from 1985 to 2017. C, countries with an urban excess BMI in 1985 that changed to a rural excess BMI in 2017. D, countries with a rural excess BMI that increased from 1985 to 2017. E, countries with a rural excess BMI that decreased from 1985 to 2017. F, countries with a rural excess BMI in 1985 that changed to an urban excess BMI in 2017.



Extended Data Fig. 5 | The relationship between mean BMI and prevalence of overweight. Overweight is defined as BMI ≥ 25 kg m⁻². Prevalence is plotted on a probit scale, which changes in an approximately linear manner as the mean changes. Each point represents an age group-

and sex-specific mean, stratified by place of residence as described in the Methods and with more than 25 participants, from data sources in the NCD-RisC database.



Extended Data Fig. 6 | Comparison of the difference between rural and urban age-standardized mean BMI in women and men aged 18 years and older in 1985 and 2017. Each point shows one country and colours indicate region.

Extended Data Table 1 | Mean BMI and percentage of the population by urban and rural place of residence

Region	Sex	Percentage of the population living in urban areas		Age-standardized mean BMI (kg m ⁻²)			
				Rural		Urban	
		1985	2017	1985	2017	1985	2017
Emerging economies							
Central Asia, Middle East and north Africa	Men	51%	63%	23.4 (22.7-24.0)	26.0 (25.8-26.3)	24.2 (23.6-24.8)	26.8 (26.6-27.0)
	Women			24.2 (23.5-24.9)	28.2 (27.9-28.4)	26.1 (25.4-26.8)	28.7 (28.4-28.9)
East and southeast Asia	Men	25%	55%	20.7 (20.4-21.0)	23.3 (23.0-23.7)	21.8 (21.5-22.1)	24.4 (24.0-24.9)
	Women			20.9 (20.5-21.3)	23.3 (22.9-23.8)	22.1 (21.7-22.4)	23.9 (23.4-24.4)
Latin America and the Caribbean	Men	68%	80%	23.0 (22.4-23.6)	25.6 (25.3-25.9)	24.4 (23.8-24.9)	26.9 (26.6-27.2)
	Women			23.1 (22.5-23.7)	27.1 (26.7-27.4)	24.6 (24.0-25.1)	27.5 (27.2-27.9)
Oceania	Men	21%	20%	22.9 (21.9-24.0)	25.7 (24.7-26.7)	25.3 (24.2-26.4)	26.6 (25.7-27.5)
	Women			23.2 (21.9-24.5)	26.2 (24.8-27.5)	25.6 (24.1-27.1)	28.4 (27.2-29.6)
South Asia	Men	24%	34%	19.0 (18.4-19.6)	21.6 (21.2-22.0)	22.0 (21.3-22.6)	22.8 (22.4-23.2)
	Women			19.0 (18.2-19.7)	21.8 (21.4-22.3)	22.2 (21.3-23.0)	23.7 (23.3-24.2)
Sub-Saharan Africa							
Sub-Saharan Africa	Men	25%	39%	20.1 (19.6-20.7)	21.7 (21.3-22.0)	21.7 (21.2-22.3)	23.3 (22.9-23.7)
	Women			20.9 (20.4-21.5)	22.7 (22.4-23.0)	23.5 (23.0-24.1)	25.9 (25.6-26.2)
High-income and other industrialized regions							
Central and eastern Europe	Men	65%	68%	25.0 (24.4-25.6)	26.7 (26.2-27.2)	25.0 (24.5-25.5)	26.7 (26.1-27.3)
	Women			26.3 (25.6-27.0)	26.7 (26.1-27.2)	26.0 (25.4-26.6)	26.2 (25.6-26.9)
High-income Asia Pacific	Men	74%	91%	22.3 (22.0-22.6)	24.1 (23.9-24.4)	22.4 (22.1-22.6)	23.9 (23.6-24.2)
	Women			22.2 (21.8-22.6)	22.7 (22.3-23.0)	22.1 (21.8-22.4)	22.1 (21.7-22.5)
High-income western countries	Men	74%	80%	25.6 (25.3-25.8)	27.8 (27.5-28.1)	25.2 (24.9-25.4)	27.6 (27.4-27.9)
	Women			25.4 (25.0-25.8)	26.9 (26.5-27.3)	24.9 (24.6-25.2)	26.9 (26.5-27.2)
World							
World	Men	41%	55%	21.1 (20.9-21.3)	23.2 (23.0-23.4)	23.7 (23.5-23.9)	25.3 (25.2-25.5)
	Women			21.5 (21.3-21.8)	23.6 (23.4-23.8)	24.2 (23.9-24.4)	25.5 (25.3-25.7)

For each region, the table shows age-standardized mean BMI for urban and rural populations and the percentage of the population living in urban areas in 1985 and 2017. See Supplementary Table 2 for a list of countries in each region. Numbers in parentheses show 95% credible intervals.

Extended Data Table 2 | Results of model validation

Validation	Sex	Data		No. of held-out observations	Percent covered	Error (kg m ⁻²) [†]				Absolute error (kg m ⁻²)			
						Median	Q1	Q3	(p*)	Median	Q1	Q3	(p*)
Test 1	Women	All		7022	98	0.03	-1.09	1.08	0.08	1.09	0.50	1.88	0.08
		Study representativeness	Community	1589	98	0.30	-0.89	1.31	0.48	1.16	0.54	1.95	0.48
			Sub-national	1197	98	-0.09	-1.30	1.25	0.03	1.29	0.61	2.04	0.03
			National	4236	97	-0.03	-1.09	0.95	0.06	1.00	0.47	1.78	0.06
		Years	1985-1999	480	96	0.00	-1.33	0.86	0.14	1.06	0.47	1.69	0.14
			2000-2017	6542	98	0.03	-1.07	1.10	0.17	1.09	0.50	1.90	0.17
	Men	All		6392	97	-0.15	-0.98	0.74	0.00	0.86	0.43	1.54	0.00
		Study representativeness	Community	1409	98	-0.15	-0.92	0.63	0.01	0.78	0.39	1.37	0.01
			Sub-national	1233	98	-0.11	-0.85	0.87	0.18	0.86	0.40	1.56	0.18
			National	3750	96	-0.16	-1.06	0.74	0.00	0.89	0.45	1.59	0.00
		Years	1985-1999	627	99	-0.24	-0.90	0.50	0.02	0.72	0.36	1.21	0.02
			2000-2017	5765	97	-0.13	-1.00	0.76	0.00	0.88	0.44	1.58	0.00
Test 2	Women	All		7680	94	-0.03	-0.67	0.58	0.29	0.62	0.28	1.19	0.29
		Study representativeness	Community	1480	88	0.10	-0.75	0.92	0.83	0.84	0.38	1.59	0.83
			Sub-national	1330	93	-0.07	-0.73	0.58	0.38	0.65	0.30	1.19	0.38
			National	4870	96	-0.05	-0.63	0.50	0.27	0.57	0.26	1.08	0.27
		Years	1985-1999	1472	94	-0.02	-0.56	0.55	0.71	0.56	0.25	1.15	0.71
			2000-2017	6208	94	-0.03	-0.69	0.58	0.21	0.64	0.29	1.20	0.21
	Men	All		6608	95	0.00	-0.54	0.51	0.23	0.52	0.24	1.01	0.23
		Study representativeness	Community	1559	93	0.01	-0.65	0.65	0.71	0.65	0.32	1.23	0.71
			Sub-national	1137	94	0.03	-0.51	0.56	0.93	0.54	0.25	1.06	0.93
			National	3912	96	-0.01	-0.51	0.45	0.19	0.48	0.21	0.91	0.19
		Years	1985-1999	1190	97	-0.04	-0.52	0.37	0.68	0.45	0.20	0.84	0.68
			2000-2017	5418	95	0.01	-0.55	0.53	0.28	0.54	0.25	1.05	0.28

Q1, first quartile; Q3, third quartile; p, p value.

[†]Estimated values minus held-out values.

*p values for model error comparisons were calculated using the non-parametric Wilcoxon signed-rank test for paired data. The p values are calculated assuming independence of the held-out observations. They should therefore be interpreted as an approximation because there is some dependence among the held-out observations, within each of the five repetitions for example.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a Confirmed

- ☒ ☐ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☒ ☐ An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☒ ☐ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☒ ☐ A description of all covariates tested
- ☒ ☐ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☒ ☐ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☐ ☒ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated
- ☐ ☒ Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection

Processing of secondary data was conducted using the statistical software R (version 3.5.1).

Data analysis

All analyses were conducted using the statistical software R (version 3.5.1). The code for national analysis of mean risk factor trends is available at www.ncdrisc.org. The code for analysis of trends in urban and rural subgroups is available from www.ncdrisc.org.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

This is a data-pooling study that brings together almost 2000 disparate data sources and uses a Bayesian hierarchical model to estimate population risk factor trends. Estimates of mean BMI by country, year, sex and place of residence (urban and rural) are available from www.ncdrisc.org. Estimates of mean BMI by

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☒ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/authors/policies/ReportingSummary-flat.pdf](https://www.nature.com/authors/policies/ReportingSummary-flat.pdf)

Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We pooled and re-analysed population-based data on height and weight in adults to estimate trends in mean BMI from 1985 to 2017 by urban and rural place of residence from 200 countries and territories, using a Bayesian hierarchical model.
Research sample	We pooled data from 2,009 population-based studies of human anthropometry conducted in 190 countries, with measurement of height and weight in over 112 million adults aged 18 years and older. Studies were representative of a national, subnational or community population.
Sampling strategy	We included data collected using a probabilistic sampling method with a defined sampling frame. We therefore included studies with simple random and complex survey designs but excluded convenience samples.
Data collection	We used data on measured height and weight to calculate body-mass index. We excluded self-reported data.
Timing	We pooled data collected from 1985 to 2017. We also included national studies for the 3 years prior to 1985 (n=17), assigning them to 1985, so that they can inform the estimates in countries with slightly earlier national data.
Data exclusions	We excluded all data sources that were solely based on self-reported weight and height without a measurement component because these data are subject to biases that vary by geography, time, age, sex and socioeconomic characteristics. Due to these variations, approaches to correcting self-reported data leave residual bias. We also excluded data sources on population subgroups whose anthropometric status may differ systematically from the general population, including: • studies that had included or excluded people based on their health status or cardiovascular risk; • studies whose participants were only ethnic minorities; • specific educational, occupational, or socioeconomic subgroups, with the exception noted below; • those recruited through health facilities, with the exception noted below; and • women aged 18-19 years in surveys which sampled only ever-married women or measured height and weight only among mothers. Our exclusion criteria were established at the initiation of the study to ensure all data were representative.
Non-participation	Our inclusion/exclusion criteria were designed to ensure participants of the surveys included were representative of the general population from which each sample was drawn.
Randomization	Our study is descriptive, and we did not carry out experiments.

Reporting for specific materials, systems and methods

Materials & experimental systems

n/a	Involved in the study
☒	☐ Unique biological materials
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology
☒	☐ Animals and other organisms
☒	☐ Human research participants

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging